

Sub-critical and Super-critical Regimes in Epidemic Models of Earthquake Aftershocks

Agnès Helmstetter

Laboratoire de Géophysique Interne et Tectonophysique, Observatoire de Grenoble, Université Joseph Fourier, France

Didier Sornette

Laboratoire de Physique de la Matière Condensée, CNRS UMR 6622 Université de Nice-Sophia Antipolis, Parc Valrose, 06108 Nice, France and Department of Earth and Space Sciences and Institute of Geophysics and Planetary Physics, University of California, Los Angeles, California 90095-1567

Abstract

We present an analytical solution and numerical tests of the epidemic-type aftershock (ETAS) model for aftershocks, which describes foreshocks, aftershocks and mainshocks on the same footing. In this model, each earthquake of magnitude m triggers aftershocks with a rate proportional to $10^{\alpha m}$. The occurrence rate of aftershocks triggered by a single mainshock decreases with the time from the mainshock according to the modified Omori law $K/(t+c)^p$ with $p = 1 + \theta$. Contrary to the usual definition, the ETAS model does not impose an aftershock to have a magnitude smaller than the mainshock. Starting with a mainshock at time $t = 0$ that triggers aftershocks according to the local Omori law, that in turn trigger their own aftershocks and so on, we study the seismicity rate of the global aftershock sequence composed of all the secondary and subsequent aftershock sequences. The effective branching parameter n , defined as the mean aftershock number triggered per event, controls the transition between a sub-critical regime $n < 1$ to a super-critical regime $n > 1$. A characteristic time t^* , function of all the ETAS parameters, marks the transition from the early time behavior to the large time behavior. In the sub-critical regime, we recover and document the crossover from an Omori exponent $1 - \theta$ for $t < t^*$ to $1 + \theta$ for $t > t^*$ found previously in [Sornette and Sornette, 1999] for a special case of the ETAS model. In the super-critical regime $n > 1$ and $\theta > 0$, we find a novel transition from an Omori decay law with exponent $1 - \theta$ for $t < t^*$ to an explosive exponential increase of the seismicity rate for $t > t^*$. The case $\theta < 0$ yields an infinite n -value. In this case, we find another characteristic time τ controlling the crossover from an Omori law with exponent $1 - |\theta|$ for $t < \tau$, similar to the local law, to an exponential increase at large times. These results can rationalize many of the stylized facts reported for aftershock and foreshock sequences, such as (i) the suggestion [Liu, 1984; Bowman, 1997] that a small p -value may be a precursor of a large earthquake, (ii) the relative seismic quiescence sometimes observed before large aftershocks, (iii) the positive correlation between b and p -values, (iv) the observation that great earthquakes are sometimes preceded by a decrease of b -value and (v) the acceleration of the seismicity preceding great earthquakes.

Introduction

It is well known that the seismicity rate increases after a large earthquake, for time period up to one hundred years [Utsu *et al.*, 1995], and distances up to several hundred km [Tajima and Kanamori, 1985; Steeples and Steeples, 1996; Kagan and Jackson, 1998; Meltzner and Wald, 1999; Dreger and Savage, 1999]. The rate of the triggered events usually decays in time as the modified Omori law $n(t) = K/(t + c)^p$, where the exponent p is found to vary between 0.3 and 2 [Davis and Frohlich, 1991; Kisslinger and Jones, 1991; Guo and Ogata, 1995; Utsu *et al.*, 1995] and is often close to 1 (see however [Kisslinger, 1993; Gross and Kisslinger, 1994] for alternative decay laws such as the stretched exponential).

These triggered events are called aftershocks if their magnitude is smaller than the first event. However, the definition of an aftershock contains unavoidably a degree of arbitrariness because the qualification of an earthquake as an aftershock requires the specification of time and space windows. In this spirit, several alternative algorithms for the definition of aftershocks have been proposed [Gardner and Knopoff, 1974; Molchan and Dmitrieva, 1992] and there is no consensus.

Aftershocks may result from several and not necessarily exclusive mechanisms (see [Harris, 2001] and references therein): pore-pressure changes due to pore-fluid flows coupled with stress variations, slow redistribution of stress by aseismic creep, rate-and-state dependent friction within faults, coupling between the viscoelastic lower crust and the brittle upper crust, stress-assisted micro-crack corrosion [Yamashita and Knopoff, 1987; Lee and Sornette, 2000], slow tectonic driving of a hierarchical geometry with avalanche relaxation dynamics [Huang *et al.*, 1998], dynamical hierarchical models with heterogeneity, feedbacks and healing [Blanter *et al.*, 1997], etc.

Since the underlying physical processes are not fully understood, the qualifying time and space windows are more based on common sense than on hard science. Particularly, there is no agreement about the duration of the aftershock sequence and the maximum distance between aftershock and mainshock. If one event occurs with a magnitude larger than the first event, it becomes the new mainshock and all preceding events are retrospectively called foreshocks. Thus, there is no way to identify foreshocks from usual aftershocks in real time. There is also no way to distinguish aftershocks from individual earthquakes [Hough

and Jones, 1997]. The aftershock magnitude distribution follows the Gutenberg-Richter distribution with similar b -value as other earthquakes [Ranalli, 1969; Knopoff *et al.*, 1982]. They have also similar rupture process. Moreover, an event can be both an aftershock of a preceding large event, and a mainshock of a following earthquake. For example, the M=6.5 Big Bear event is usually considered as an aftershock of the M=7.3 Landers event, and has clearly triggered its own aftershock sequence. One can trace the difficulty of the problem from the long-range nature of the interactions between faults in space and time resulting in a complex self-organized crust.

In view of the difficulties in classifying sometimes an earthquake as a foreshock, a mainshock or an aftershock, it is natural to investigate models in which this distinction is removed and to study their possible observable consequences. In this spirit, the epidemic type aftershock (ETAS) model introduced by Kagan and Knopoff [1981, 1987] and Ogata [1988] provides a tool for understanding the temporal clustering of the seismic activity without distinguishing between aftershocks, foreshocks and mainshock events. The ETAS model is a generalization of the modified Omori law, which takes into account the secondary aftershocks sequences triggered by all events. In this model, all earthquakes are simultaneously mainshocks, aftershocks and possibly foreshocks. An observed “aftershock” sequence is in the ETAS model the result of the activity of all events triggering events triggering themselves other events, and so on, taken together. The ETAS model aims at modeling complex aftershocks sequences and global seismic activity. The seismicity rate is given by the superposition of aftershock sequences of all events. Each earthquake of magnitude m triggers aftershock with a rate proportional to $10^{\alpha m}$ with the same coefficient α for all earthquakes. The occurrence rate of aftershocks decreases with the time from the mainshock according to the modified Omori law $K/(t + c)^p$. The background seismicity rate is modeled by a stationary Poisson process with a constant occurrence rate μ . Contrary to the usual definition, the ETAS model does not impose an aftershock to have a magnitude smaller than the mainshock. This way, the same law describes both foreshocks, aftershocks and mainshocks. This model has been used to give short-term probabilistic forecast of seismic activity [Kagan and Knopoff, 1987; Kagan and Jackson, 2000; Console and Murru, 2001], and to describe the temporal and spatial clustering of seismic activity [Ogata, 1988, 1989, 1992, 1999, 2001;

Kagan, 1991; Felzer et al, 2001]. Although the elementary results on the stability of the process have been known for many years [*Kagan, 1991*], no attempt has been made to study this model analytically in order to characterize its different regimes and obtain a deeper understanding of the combined interplay between the model parameters (b , α , p , K , c and μ) on the seismic activity. We stress below the contrast between previous works in the mathematical statistical literature and our results.

It should be noted that the ETAS model suffers from an important defect: it is fundamentally a “branching” model [*Harris, 1963; Vere-Jones, 1977*], with no “loops”. What this means is that an event has a unique “mother-mainshock” and not several. In the real case, we can expect that some events may be triggered by the combined loading and action at distance in time and space of several previous earthquakes. Hence, events should have several “mothers” in general. This neglecting of “loops” is known in statistical physics as a “mean-field” approximation and allows us to simplify the analysis while still keeping the essential physics in a qualitative way, even if the details may not be precisely recovered quantitatively.

Sornette and Sornette [1999] studied analytically a particular case of the ETAS model, in which the aftershock number does not depend on the mainshock magnitude, *i.e.*, for $\alpha = 0$. Starting with one event at time $t = 0$ and considering that each earthquake generates an aftershock sequence with a “local” Omori exponent $p = 1 + \theta$, where θ is a positive constant, they studied the decay law of the “global” aftershock sequence, composed of all secondary aftershock sequences. They found that the global aftershock rate decays according to an Omori law with an exponent $p = 1 - \theta$, smaller than the local one, up to a characteristic time t^* , and then recovers the local Omori exponent $p = 1 + \theta$ for time larger than t^* .

Here, we generalize their analysis in the more general case $\alpha > 0$ of the ETAS model, which includes a realistic magnitude distribution. We study the decay law of the global aftershock sequence as a function of the model parameters (local Omori law parameters and magnitude distribution). In addition to giving more complete analytical results, we present numerical simulations that test these predictions. We also generalize the investigation and analysis into the “super-critical” regime. Indeed, depending on the branching ratio n , defined as the mean aftershock number triggered per event, and on the sign of θ , three different regimes for the seismic rate $N(t)$ are found:

1. For $n < 1$ (sub-critical regime), we recover the results of [*Sornette and Sornette, 1999*], *i.e.* we find a crossover from an Omori exponent $p = 1 - \theta$ for $t < t^*$ to $p = 1 + \theta$ for $t > t^*$.
2. For $n > 1$ and $\theta > 0$ (super-critical regime), we find a transition from an Omori decay law with exponent $p = 1 - \theta$ to an explosive exponential increase of the seismicity rate.
3. In the case $\theta < 0$, we find a transition from an Omori law with exponent $1 - |\theta|$ similar to the local law, to an exponential increase at large times, with a crossover time τ different from the characteristic time t^* found in the case $\theta > 0$.

As we show below, these results can rationalize many properties of aftershock and foreshock sequences.

The model

We assume that a given event (the “mother”) of magnitude $m_i \geq m_0$ occurring at time t_i gives birth to other events (“daughters”) in the time interval between t and $t + dt$ at the rate

$$\phi_{m_i}(t - t_i) = \frac{K 10^{\alpha(m_i - m_0)}}{(t - t_i + c)^{1+\theta}} H(t - t_i) H(m_i - m_0), \quad (1)$$

where H is the Heaviside function: $H(t - t_i) = 0$ for $t < t_i$ and 1 otherwise, m_0 is a lower bound magnitude below which no daughter is triggered.

This temporal power law decay is Omori’s law for the rate of aftershocks following a main shock, albeit with the modification that we do not specify that aftershocks (daughter earthquakes) have to be smaller than the triggering event (mother earthquake). The exponential term $10^{\alpha(m - m_0)}$ describes the fact that the larger the magnitude m of the mother event, the larger is the number of daughters. The exponent $p = 1 + \theta$ of the “local” Omori’s law has no reason a priori to be the same as the one measured macroscopically which is usually found between 0.8 and 1.2 with an often quoted median value 1. This is in fact the question we address: assuming the form (1) for the “local” Omori’s law, is the global Omori’s law still a power law and, if yes, how does its exponent depend on p ? What are the possible regimes of aftershocks as a function of the parameters of the model?

This model can be extended to describe the spatio-temporal distribution of seismic activity. Following

Kagan and Knopoff [1981], we can introduce a spatial dependence in (1) of the form

$$\phi_{m_i}(t-t_i, \vec{r}-\vec{r}_i) = \frac{K}{(t-t_i+c)^{1+\theta}} 10^{\alpha(m_i-m_0)} \rho(\vec{r}-\vec{r}_i) H(t-t_i) H(m_i-m_0), \quad (2)$$

where $\rho(\vec{r}-\vec{r}_i)$ describes the probability distribution for an earthquake occurring at position \vec{r}_i to trigger an event at position \vec{r} . This term takes into account the spatial dependence of the stress induced by an earthquake, and enable us to model the spatial distribution of aftershocks clustered close to the mainshock. In this paper, we restrict our analysis to the temporal ETAS model without spatial dependence because we are mainly interested in describing the temporal evolution of seismic activity. The complete model with both spatial and temporal dependence (2) has been studied in [*Helmstetter and Sornette, 2002*] to derive the joint probability distribution of the times and locations of aftershocks including the whole cascade of secondary aftershocks. When integrating the rate of aftershocks calculated for the spatio-temporal ETAS model over the whole space, we recover the results given in this paper for the temporal ETAS model. Therefore, the results given here for the temporal ETAS model can be compared with real aftershock sequences when using all aftershocks whatever their distance from the mainshock.

The model (1) is a branching process because each daughter has only one mother and not several, as shown in Figure 1. As we said in the introduction, this “mean-field” assumption simplifies considerably the complexity of the process and allows for an analytical solution that we shall derive in the sequel. The key parameter is the average number n of daughter-earthquakes created per mother-event. Assuming that the distribution $P(m)$ of earthquake sizes expressed in magnitudes m follows the Gutenberg-Richter distribution $P(m) = b \ln(10) 10^{-b(m-m_0)}$, the integral of $\phi_m(t)$ over time and over all magnitudes $m \geq m_0$ gives

$$n \equiv \int_0^{+\infty} dt \int_{m_0}^{+\infty} dm P(m) \phi_m(t) = n_0 \int_0^{+\infty} \frac{dt}{(t+1)^{1+\theta}}, \quad (3)$$

where

$$n_0 \equiv \frac{K}{c^\theta} \frac{b}{b-\alpha}, \quad (4)$$

which is finite for $b > \alpha$. Three cases are analyzed below: $n < 1$, $n = 1$ and $n > 1$. The case $n = 1$ corresponds to an average conservation of the number of events and can be associated with a brittle elas-

tic crust without dissipation. The “dissipative” case $n < 1$ can be interpreted as corresponding to a crust possessing a visco-elastic component and/or a partial coupling with a lower ductile layer, such that a part of the energy is released aseismically. The case $n > 1$ corresponds to a process in which an earthquake sequence triggers an in-flow of energy from surrounding regions that may lead to a local self-exciting amplification. It can also correspond to a coupling with other non-mechanical modes of energy storage, such as proposed in [*Sornette, 2000b; Viljoen et al., 2002*] which can be triggered by an event and feed the ensuing earthquake sequence for a while. Of course, the super-critical process can only be transient and has to cross-over to another regime.

The case $b < \alpha$ requires a special attention. In absence of truncation or cut-off, it leads to a finite-time singularity due to the interplay between long-memory and extreme fluctuations [*Sornette and Helmstetter, 2001*]. However, it is more common to introduce a truncation or roll-off of the Gutenberg-Richter law at an upper magnitude. We can for example use a Gamma distribution of energies, which is a power-law distribution tapered by an exponential tail. In this case, the branching ratio has been calculated by *Kagan* [1991] and is given by the approximate analytical expression valid for a corner magnitude m_c significantly larger than m_0 ,

$$n_0 = \frac{K}{c^\theta} \frac{b}{b-\alpha} \frac{10^{b(m_c-m_0)} - 10^{\alpha(m_c-m_0)}}{10^{b(m_c-m_0)} - 1}. \quad (5)$$

For a corner magnitude $m_c \gg m_0$, and for $\alpha < b$, we recover the expression (4) for n_0 obtained for the Gutenberg-Richter distribution without roll-off.

Note that n is defined as the average over all mainshock magnitudes of the mean number of events triggered by a mainshock. It is thus grossly misleading to think of the branching ratio as giving the number of daughters to a given earthquake, because this number is extremely sensitive to the specific value of its magnitude. Indeed, the number of aftershocks to a given mainshock increases exponentially with the mainshock magnitude as given by (1), so that large earthquakes will have many more aftershocks than small earthquakes. From (1) and (3), we can calculate the mean number of aftershocks $N(M)$ triggered directly by a mainshock of magnitude M

$$N(M) = n \frac{(b-\alpha)}{b} 10^{\alpha(M-m_0)}. \quad (6)$$

As an example, take $\alpha = 0.8$, $b = 1$, $m_0 = 0$ and $n = 1$. Then, a mainshock of magnitude $M = 7$ will

have on average 80000 direct aftershocks, compared to only 2000 direct aftershocks for an earthquake of magnitude $M = 5$ and less than 0.2 aftershocks for an earthquake of magnitude $M = 0$.

When $\theta > 0$, $\int_0^\infty \frac{dt}{(t+1)^{1+\theta}} = 1/\theta$ and the branching ratio $n = n_0/\theta$ is finite. In this regime, n is an increasing function of the rate K and a decreasing function of θ , c and $b - \alpha$.

Even for $b > \alpha$ and $\theta > 0$, the average number of daughters per mother can be larger than one: $n > 1$. This regime corresponds to the super-critical regime of branching processes [Harris, 1963; Sornette, 2000a] in which the total number of events grows on average exponentially with time. If $n < 1$, there is less than one earthquake triggered per earthquake on average. This is the sub-critical regime in which the number of events following the first main shock decays eventually to zero. The critical case $n = 1$ is at the borderline between the two regimes. In this case, there is exactly one earthquake on average triggered per earthquake and the process is exactly at the critical point between death on the long run and exponential proliferation.

There is another scenario, occurring for $\theta \leq 0$, in which the seismicity blows up exponentially with time. In this case, the integral $\int_0^\infty \frac{dt}{(t+1)^{1+\theta}}$ becomes unbounded. In principle, n becomes infinite: this does not invalidate the ETAS model per se. It only reflects the fact that the calculation of an average number of daughters per mother has become meaningless because of the anomalously slow decay of the kernel $\phi(t)$. This mechanism is reminiscent of that leading to anomalous diffusion and to aging in quenched random systems and spinglasses (see [Sornette, 2000a] for an introduction). As in these systems, any estimation of the averages depend on the time scale of study: due to the extremely slow decay of $\phi(t)$, the number of daughters created beyond any time t far exceeds the number of daughters created up to time t . Notwithstanding the decay, its cumulative effect creates this dominance of the far future. This regime is the opposite of the situation where $\theta > 0$ where most of the daughters are created at relatively early times. Since the number of daughters born up to time t is an unbounded increasing function of t , it is intuitively appealing, as we show in the appendix, that this regime should be similar to the super-critical regime $n > 1$ discussed above in the case $\theta > 0$.

Until now, we have discussed three issues related to the convergence of the ETAS sequences: (i) the

condition $\theta > 0$ ensures convergence at large times; (ii) the convergence at short times is obtained by the introduction of the regularization constant c in the generalized Omori's law; (iii) the condition $\alpha < b$ is a necessary condition for the finiteness of the number of daughters. Finally, we should stress the role of the "ultra-violet" cut-off m_0 on the magnitudes. In the ETAS model, only earthquakes of magnitude $m \geq m_0$ are allowed to give birth to aftershocks, while events of smaller magnitudes are lost for the epidemic dynamics. If such a cut-off is not introduced and no cut-off is put on the Gutenberg-Richter toward small magnitudes, the dynamics becomes completely dominated by the swarms of very tiny earthquakes, which individually has very low probability to generate aftershocks but become so numerous that their collective effect becomes overwhelming in the dynamics. We would thus have the unphysical situation in which a magnitude 7 or 8 earthquake may be triggered by tiny earthquakes of magnitudes -2 or less. We stress that the introduction of such a cut-off m_0 is a simple way to prevent such a situation to occur, but it does not mean that small earthquakes of magnitude below m_0 do not have their own aftershocks. It only means that such small earthquakes create aftershocks that can not participate in the epidemic process leading to significantly larger earthquakes; these small earthquakes live their separate life. This is why they are not registered by the ETAS model. This formulation is of course only an end-member of many possible regularization procedures, which are well-known to be an ubiquitous requisite in mechanical models of rupture. An improvement of the ETAS model would be for instance to replace this abrupt cut-off m_0 by introducing a roll-off in the Gutenberg-Richter law for the aftershocks with a characteristic corner magnitude decreasing with the magnitude of the mother earthquake. This and other schemes will not be explored here, as we want to analyze the simplest version possible.

We now describe briefly the connection with previous works in the mathematical statistics literature. As we said above, the model (1) belongs to the general class of branching models [Moyal, 1962; Harris, 1963]. The elementary results on the stability of the process, such as the condition $n < 1$, have been known for many years, and go back to the origin of the ETAS model as a special case (for discrete magnitudes) or extension (for continuous magnitudes) of the class of "mutually exciting point processes" introduced in [Hawkes, 1971; 1972; Hawkes

and Adamopoulos, 1973]. A convenient mathematical overview is in Chapter 5 of Vere-Jones and Daley [1988], especially Example 5.5(a) and associated exercises 5.5.2-5.5.6. For the ETAS model, the equations governing the probability generating functional, the probability of extinction within a given number of generations, the expectation measure for the total population, the second factorial moment (related to the covariance of the population) and their Fourier transform can be derived as special cases of results summarized there. In particular, the process initiated with a single event at the origin corresponds to the total progeny process for a general branching process model with time-magnitude state space and a single ancestor at time $t = 0$; Exercise 5.5.6 gives the equations of the above cited variables for the case of fixed magnitudes (i.e., $\alpha = 0$). This direct probabilistic analysis in terms of generating functions effectively replaces the Wiener-Hopf theory in the present paper and mentioned also in [Hawkes, 1971; 1972; Hawkes and Oakes, 1974]. However, there is not explicit solutions given to these equations and there is no discussion of the change of regime from an effective Omori's law $1/t^{1-\theta}$ at early times to $1/t^{1+\theta}$ at long times, nor mention of the interesting super-critical case, as done in the present work.

Hawkes [1971; 1972] and Hawkes and Adamopoulos [1973] use what is in effect an ETAS model with an exponential “bare” Omori's law rather than the power law $1/(t+c)^{1+\theta}$ defined in (1). Hawkes and Adamopoulos [1973] use it in an early study of earthquake data. The introduction of magnitudes is similar to the introduction of a marked process associated with a single point process [Hawkes, 1972]; however, the impact of magnitudes on the seismicity rate is assumed to be linear in [Hawkes, 1972] while it is multiplicative in the ETAS model. Our derivation presented in the appendix of the solution of the ETAS model for the mean rate of earthquakes in terms of its Laplace transform recovers previous results. For instance, equation (17) in [Hawkes and Oakes, 1974] is the same as our equation (A6) in our Appendix (up to a factor β stemming from taking the cumulative number in [Hawkes and Oakes, 1974]). The key factor $Q(\beta)$ in (A7) corresponds to the quantity $G_1(0)$ in equation (5) of [Hawkes, 1972]. The link between Hawkes' “mutually exciting point processes” and branching processes was made explicit in [Hawkes and Oakes, 1974].

Some average properties of the ETAS model have been derived in the Master thesis of P.A. Ramse-

laar (1990). Specifically, using the theory of Markov processes applied to branching processes, Ramselaar (1990) proves that, in the supercritical regime $n > 1$ (where n is the average branching ratio defined in (5)), the average number of aftershocks stemming from a common ancestor grows exponentially as $\sim e^{t/t^*}$ where t^* is the solution of $nR(c/t^*) = 1$ and the function R is defined in (A9). The solution of this equation $nR(c/t^*) = 1$ for t^* is the same as our t^* given by (12) and the exponential growth of Ramselaar is therefore the same as our result (17). We add on this asymptotic result, which is valid only at large times, by exhibiting the solution for the aftershock decay at early times. In addition, contrary to the incorrect claim of Ramselaar (1990) that “the Ogata earthquake process is critical or supercritical but is never subcritical,” we demonstrate that the subcritical regime exhibits a rich phenomenology.

Analytical solution

We analyze the case where there is an origin of time $t = 0$ at which we start recording the rate of earthquakes, assuming that the largest earthquake of all has just occurred at $t = 0$ and somehow reset the clock. In the following calculation, we will forget about the effect of events preceding the one at $t = 0$ and count aftershocks that are created only by this main shock.

Let us call $N_m(t)$ the rate of seismicity at time t and at magnitude m , that is, $N_m(t)dtdm$ is the number of events in the time/magnitude interval $dt \times dm$. We define its expectation $\lambda_m(t)dtdm \equiv E[N_m(t) dt dm]$, as the mean number of earthquakes occurring between t and $t + dt$ of magnitude between m and $m + dm$. $\lambda_m(t)$ is the solution of a self-consistency equation that formalizes mathematically the following process: an earthquake may trigger aftershocks; these aftershocks may trigger their own aftershocks, and so on. The rate of seismicity at a given time t is the result of this cascade process. The self-consistency equation that sums up this cascade reads

$$\begin{aligned} \lambda_m(t) \equiv E[N_m(t)] &= E \left[\int_{m_0}^{\infty} dm' \int_{-\infty}^t d\tau \phi_{m'}(t-\tau) P(m) N_{m'}(\tau) \right] \\ &= \int_{m_0}^{\infty} dm' \int_{-\infty}^t d\tau \phi_{m'}(t-\tau) E[N_{m'}(\tau)] \\ &= \int_{m_0}^{\infty} dm' \int_{-\infty}^t d\tau \phi_{m'}(t-\tau) \lambda_{m'}(\tau). \end{aligned}$$

If there is an external source $S(t, m)$, it should be added to the right-hand-side of (9).

The mean instantaneous rate $\lambda_m(t)$ at time t is the sum over all induced rates from all earthquakes of all possible magnitudes that occurred at all previous times. The rate of events at time t induced per earthquake that occurred at an earlier time τ with magnitude m' is equal to $\phi_{m'}(t - \tau)$. The term $P(m)$ is the probability that an event triggered by an earthquake of magnitude m' is of magnitude m . We assume that this probability is independent of the magnitude of the mother-earthquake and is nothing but the Gutenberg-Richter law. This hypothesis can be easily relaxed if needed and $P(m)$ can be generalized into $P(m|m')$ giving the probability that a daughter-earthquake is of magnitude m conditioned on the value m' of the magnitude of the mother-earthquake. However, we do not pursue here this possibility as this hypothesis seems well-founded empirically [Ranalli, 1969; Knopoff et al., 1982]. The term $S(t, m)$ is an external source which is determined by the physical process. We consider the case where a great earthquake occurs at the origin of time $t = 0$ with magnitude M . In this case, the external source term is

$$S(t, m) = \delta(t) \delta(m - M) , \quad (10)$$

where δ is the Dirac distribution. Other arbitrary source functions can be chosen.

By construction of the kernel (1), the integrals over the magnitudes and over time factorize, which implies that the solution for $\lambda_m(t)$ can be searched as

$$\lambda_m(t) = P(m)\lambda(t) , \quad (11)$$

which makes explicit the separation of the variables magnitude and time. We stress that (11) is not an assumption: it is the structure of the solution based on the assumption made at the level of equations (1) and (9). Therefore, the ETAS model assumes that the time response and the magnitude response are independent. In reality and more generality, we can envision that the rate of activation of new earthquakes will depend on 1) the magnitude of the “mother” (which the ETAS model takes into account multiplicatively in (1)), 2) on the magnitude of the daughter (which is neglected in the ETAS model) and 3) on the time since the mother was born. Rather than having a very general kernel combining these three parameters nonlinearly, equations (1) and (9) are based on an hypothesis of independence between these different factors that allows us to factorize them, leading to (11).

The problem is then to determine the functional form of $\lambda(t)$, assuming that ϕ is given by (1). The in-

tegral equation (9) is a Wiener-Hopf integral equation [Feller, 1971]. It is well-known [Feller, 1971; Morse and Feshbach, 1953] that, if $\phi(\tau)$ decays no slower than an exponential, then $\lambda(t)$ has an exponential tail $\lambda(t) \sim \exp[-rt]$ for large t with r solution of $\int \phi(x) \exp[rx] dx = 1$. This result implies that a global Omori’s law cannot be obtained by the epidemic ETAS branching model with, for instance, local exponential relaxation rates. In the present case, $\phi(\tau)$ decays much slower than an exponential and a different analysis is called for that we now present. The solution of (9) is derived in the Appendix and is summarized in the following sections. For the sequel, it is useful to define the characteristic time

$$t^* \equiv c \left(\frac{n \Gamma(1 - \theta)}{|1 - n|} \right)^{\frac{1}{\theta}} , \quad (12)$$

where $\Gamma(x)$ is the Gamma function: $\Gamma(z) = \int_0^\infty du u^{z-1} e^{-u}$ which is nothing but $(z - 1)!$ for positive integers z .

The sub-critical regime $n < 1$ and $\theta > 0$

An approximation is made in the analytical solution so that the results presented below are only valid for $t \gg c$.

We define the parameter S_0 that describes the external source term

$$S_0 = \frac{(b - \alpha)}{b} 10^{\alpha(M - m_0)} . \quad (13)$$

Two cases must be distinguished.

- For $c \ll t \ll t^*$, we get

$$\lambda_{t < t^*}(t) \sim \frac{S_0}{\Gamma(\theta)(1 - n)} \frac{t^{*-\theta}}{t^{1-\theta}} \quad \text{for } c \ll t \ll t^* . \quad (14)$$

- For $t \gg t^*$, we obtain

$$\lambda_{t > t^*}(t) \sim \frac{S_0}{\Gamma(\theta)(1 - n)} \frac{t^{*\theta}}{t^{1+\theta}} \quad \text{for } t \gg t^* . \quad (15)$$

We verify the self-consistency of the two solutions $\lambda_{t > t^*}(t)$ and $\lambda_{t < t^*}(t)$ by checking that $\lambda_{t > t^*}(t^*) = \lambda_{t < t^*}(t^*)$. In other words, t^* is indeed the transition time at which the “short-time” regime $\lambda_{t < t^*}(t)$ crosses over to the “long-time” regime $\lambda_{t > t^*}(t)$.

The full expression of $\lambda(t)$ valid at all times $t \gg c$ is given by

$$\lambda(t) = \frac{S_0}{1 - n} \frac{t^{*-\theta}}{t^{1-\theta}} \sum_{k=0}^{\infty} (-1)^k \frac{(t/t^*)^{k\theta}}{\Gamma((k + 1)\theta)} \quad (16)$$

Expression (16) provides the solution that describes the cross-over from the $1/t^{1-\theta}$ Omori's law (14) at early times to the $1/t^{1+\theta}$ Omori's law (15) at large times. The series $\sum_{k=0}^{\infty} (-1)^k \frac{(t/t^*)^{k\theta}}{\Gamma((k+1)\theta)}$ is a series representation of the special Fox function [Glöckle and Nonnenmacher, 1993] (see the Appendix for details).

The ETAS model has been simulated numerically using the algorithm described in [Ogata, 1998, 1999]. Starting with a large event of magnitude M at time $t = 0$, events are then simulated sequentially. After each event, we calculate the conditional intensity $\lambda(t)$ defined by

$$\lambda(t) = \sum_{t_i \leq t} \frac{K 10^{\alpha(m_i - m_0)}}{(t - t_i + c)^{1+\theta}}$$

where t is the time of the last event and t_i and m_i are the times and magnitudes of all preceding events that occurred at time $t_i \leq t$. The time of the following event is then determined according to the non-stationary Poisson process of conditional intensity $\lambda(t)$, and its magnitude is chosen in a Gutenberg-Richter distribution with parameter b . These simulations are compared to the theoretical predictions in Figure 2, which shows the aftershock seismic rate $\lambda(t)$ in the sub-critical regime triggered by a main event of $M = 6.8$, for the parameters $K = 0.024$ (constant in (1)), the threshold $m_0 = 0$ for aftershock triggering, $c = 0.001$, $\alpha = 0.5$, a b -value $b = 1.0$ and $\theta = 0.2$ (corresponding to a local Omori's exponent $p = 1.2$). These parameters lead to a branching ratio $n = 0.95$ (equation (3)) and a characteristic cross-over time $t^* = 4500$ (equation (12)). The noisy black line represents the seismicity rate obtained for the synthetic catalog. The local Omori law with exponent $p = 1 + \theta = 1.2$ is shown for reference as the dotted line. The analytical solution (16) is shown as the thick line. The two dashed lines represent the approximation solutions (14) for $t < t^*$ and (15) for $t > t^*$.

The super-critical regime $n > 1$ and $\theta > 0$

From the definition of the branching ETAS model for $n > 1$, it is clear that the number of events $\lambda(t)$ blows up exponentially for large times as $n - 1$ to a power proportional to the number t of generations. We shall show below that the rate of the exponential growth can be calculated explicitly, which yields $\lambda(t) \sim e^{t/t^*}$, where t^* has been defined in (12). However, there is an interesting early and intermediate time regime in the situation where a great event of magnitude M has just occurred at $t = 0$. In this

case, the total seismicity is the result of two competing effects: (1) the total seismicity tends to decay according to the Omori's law governing the rate of daughter-earthquakes triggered by the great event; (2) since each daughter may in turn trigger grand-daughters, grand-daughters may trigger grand-grand daughters and so on with a number $n > 1$ of children per parent, the induced seismicity will eventually blow up exponentially. However, before blowing up, one can expect that seismicity will first decay because it is mainly controlled by the large rate $\sim 10^{\alpha(M-m_0)}$ directly induced by the great earthquake which decays according to its "local" Omori's law. This decay will be progressively perturbed by the proliferation of daughters of daughters of ... and will cross-over to the explosive exponential regime.

At early times $c \ll t \ll t^*$, the early decay rate of aftershocks is the same $\approx (S_0/\Gamma(\theta)(n-1)) (t^{*-\theta}/t^{1-\theta})$ as for the sub-critical regime (14) (see the Appendix). However, as time increases, the Appendix shows that the decay of aftershock activity can be represented as a power law with an effective apparent exponent $\theta_{\text{app}} > \theta$ increasing progressively with time. The seismic rate will thus decay approximately as $\sim 1/t^{1-\theta_{\text{app}}(t)}$. Quantitatively, the large time behavior is (see the Appendix)

$$\lambda(t) \sim \frac{S_0}{(n-1)t^*\theta} e^{t/t^*} \quad (17)$$

exhibiting an exponential growth at large times. Expression (12) shows that $1/t^* \sim |1-n|^{1/\theta}$. Thus, as expected, the exponential growth disappears as $n \rightarrow 1^+$.

The full expression of $\lambda(t)$ valid at times $t \gg c$ is

$$\lambda(t) = \frac{S_0}{(n-1)} \frac{t^{*-\theta}}{t^{1-\theta}} \sum_{k=0}^{\infty} \frac{(t/t^*)^{k\theta}}{\Gamma((k+1)\theta)} \quad (18)$$

Expression (18) provides the solution that describes the cross-over from the $1/t^{1-\theta}$ Omori's law at early times (14) to the exponential growth (17) at large times.

Figure 3 tests these predictions by comparing them with direct numerical simulation of the ETAS model, in the case of a main shock of magnitude $M = 6$. The parameters of the synthetic catalog are $K = 0.024$ (constant in (1)), the threshold $m_0 = 0$ for aftershock triggering, $c = 0.001$, $\alpha = 0.5$, a b -value $b = 0.75$ and $\theta = 0.2$ (corresponding to a local Omori's exponent $p = 1.2$). These parameters lead to a branching ratio $n = 1.43$ (equation (3)) and a characteristic cross-over time $t^* = 0.85$ (equation (12)). The noisy black

line represents the seismicity rate obtained for the synthetic catalog. The local Omori law with exponent $p = 1 + \theta = 1.2$ is shown for reference as the dotted line. The analytical solution (18) is shown as the thick line. The two dashed lines correspond to the approximative analytical solutions (14) and (17). At early times $c < t < t^*$, the decay of $N(t)$ is initially close to the prediction (14). For $t > t^*$, we observe that the analytical equation (18) is very close to the exponential solution (17).

Case $\theta < 0$ corresponding to a local Omori's law exponent $p < 1$

We have already remarked that, in this case, the integral $\int_0^\infty \frac{dt}{(t+1)^{1+\theta}}$ in the definition (3) of the branching ratio n becomes unbounded: the number of daughters created beyond any time t far exceeds the number of daughters created up to time t .

The appendix shows that the general equation (9) still holds and the general derivation starting with (A1) up to (A6) still applies.

Similarly to the super-critical case $n > 1$ of the regime $\theta > 0$, we find a crossover from a power-law decay at early times to an exponential increase of the seismicity rate at large times. The characteristic time τ that marks the transition between these two regimes is given by

$$\tau = c \left(\frac{n_0 \Gamma(|\theta|)}{1 + \frac{n_0}{|\theta|}} \right)^{-\frac{1}{|\theta|}}. \quad (19)$$

In contrast with the case $\theta > 0$, the early time behavior (*i.e.*, $c \ll t \ll \tau$) of the global decay law in the case $\theta < 0$ is similar to the local Omori law :

$$\lambda(t) = \frac{S_0}{(1 + \frac{n_0}{|\theta|}) \Gamma(|\theta|)} \frac{\tau^{-|\theta|}}{t^{1-|\theta|}} \quad (20)$$

Similarly to the super-critical case $n > 1$ of the regime $\theta > 0$, the long time dependence of the regime $\theta < 0$ is controlled by a simple pole $1/\tau$ leading to a long-time seismicity growing exponentially

$$\lambda(t) = \frac{S_0}{(1 + \frac{n_0}{|\theta|}) \tau |\theta|} e^{t/\tau} \quad (21)$$

This result is in agreement with the fact that the number of daughters born up to time t is an unbounded increasing function of t , and we should thus recover a regime similar to the super-critical case of $\theta > 0$.

The full expression of $\lambda(t)$ valid at times $t > c$ is

$$\lambda(t) = \frac{S_0}{(1 + \frac{n_0}{|\theta|})} \frac{1}{t} \sum_{k=1}^{\infty} \frac{(t/\tau)^{k|\theta|}}{\Gamma(k|\theta|)} \quad (22)$$

Expression (22) provides the solution that describes the cross-over from the local Omori law $1/t^{1-|\theta|}$ at early times to the exponential growth at large times.

Figure 4 compares these predictions to a direct numerical simulation of the ETAS model, in the case of a main shock of magnitude $M = 7$. The parameters of the synthetic catalog are $K = 0.02$, $m_0 = 0$, $c = 0.01$, $\alpha = 0.5$, $b = 1$ and $\theta = -0.1$ (corresponding to a local Omori's exponent $p = 0.9$). These parameters lead to a characteristic cross-over time $\tau = 10^5$ (equation (19)). The noisy black line represents the seismicity rate obtained for the synthetic catalog. The local Omori law with exponent $p = 1 + \theta = 0.9$ is shown for reference as the dotted line. The analytical solution (22) is shown as the thick line. The two dashed lines correspond to the approximative analytical solutions (20) and (21). At early times $c < t < \tau$, the decay of $\lambda(t)$ is initially close to the prediction (20). For $t > \tau$, we observe that the analytical equation (22) is very close to the exponential solution (21).

Discussion

Assuming that each event triggers aftershock sequences according to the local Omori law with exponent $1 + \theta$, we have shown that the decay law of the global aftershock sequence is different from the local one. Depending on the branching ratio n , which is a function of all ETAS parameters, we find two different regimes, the sub-critical regime for $n < 1$ and the super-critical regime for $n > 1$ and $\theta > 0$. For the two regimes in the case $\theta > 0$, a characteristic time t^* , function of c , n and θ , appears in the global decay law $\lambda(t)$ and marks the transition between the early time behavior and the large time behavior. In the sub-critical regime ($n < 1$), the global decay law is composed of two power laws. At early times ($t < t^*$), $\lambda(t)$ decays like $t^{-1+\theta}$. At large times ($t > t^*$) the global decay law recovers the local law $N(t) \sim t^{-1-\theta}$. In the super-critical regime ($n > 1$ and $\theta > 0$), the early times decay law is similar to that of the sub-critical regime, and the seismicity rate increases exponentially for large times. The case $\theta < 0$ leads to an infinite n -value, due to the slow decay with time of the local Omori law. In this case, we find a transition from an Omori law with exponent $1 - |\theta|$ similar to the local law, to an exponential increase at large

times, with a crossover time τ different from the characteristic time t^* found in the case $\theta > 0$. Thus, the Omori law is only an approximation of the global decay law valid for some time periods and parameter values. The value of the local Omori exponent $p = 1$ is the only one for which the local and the global decay rate are similar, and are both power-laws without any characteristic time. For small n , t^* is very small so that in real data we should observe only the behavior $t > t^*$ characteristic of large times. The global decay law then appears similar to the local Omori law. On the contrary, for n close to 1, t^* is very large by comparison with the time period available in real data, and we should observe only the power-law behavior $\lambda(t) \sim t^{-1+\theta}$ characteristic of early times, with a global p -value smaller than the local one. Changing n thus provides an important source of variability of the exponent p .

Estimation of n and t^* in earthquake data

In real earthquake data, it is possible to evaluate the branching value n in order to determine if the seismic activity is either in the sub- or the super-critical regime. The values of n and t^* can be evaluated from equations (3) and (12) as a function of the ETAS parameters b , $p = 1 + \theta$, c , K and α . The parameters of the ETAS model and their standard error can be inverted from seismicity data (time and magnitudes of each event) using a maximum likelihood method [Ogata, 1988]. We now discuss the range of the different parameters obtained from such inversion procedure.

- The parameter α is found to vary between 0.35 to 1.7, and is often close to 0.5 [Ogata, 1989, 1992; Guo and Ogata, 1997]. An α -value of 0.5 means that a mainshock of magnitude M will have on average 10 times more aftershocks than a mainshock of magnitude $M - 2$, independently of M . Note that our definition of α is slightly different from that used by Ogata and we have divided his α -values by $\ln(10)$ to compare with our definition.

For some seismicity sequences, Ogata [1989, 1992] and Guo and Ogata [1997] found $\alpha > b$. According to (3), this leads to an infinite n -value if we use a Gutenberg-Richter magnitude distribution. As we said, a truncation of the magnitude distribution is needed to obtain a physically meaningful finite n -value because the seismicity rate is controlled by the largest

events.

A large α -value can be associated with seismic activity called “swarms”, while a small α -value is observed for aftershock sequences with a single mainshock and no significant secondary aftershock sequences [Ogata, 1992, 2001].

- The parameter c is usually found to be of the order of one hour [Utsu et al., 1995]. In practice, the evaluation of c is hindered by the incompleteness of earthquake catalogs just after the occurrence of the mainshock, due to overlapping aftershocks on the seismograms. A large c is often an artifact of a change of the detection threshold. Notwithstanding these limitations, well-determined non-zero c -value have been obtained for some aftershocks sequences [Utsu et al., 1995]. Note that a non-zero c is required for the aftershocks rate to be finite just at the time of the mainshock.
- The “local” p -value, equal to $1 + \theta$, describes the decay law of the aftershock sequence triggered by a single earthquake. The local Omori law is the law $\phi(t)$ obtained by inverting the ETAS model on the data. The “global” p -value describes the decay law of the whole aftershock sequence, composed of all secondary aftershocks triggered by each aftershock. The global Omori law is the law $\phi(t)$ fitted directly on the data, without taking into account the hierarchical structure of branching of the ETAS model. We have shown that the Omori law is only an approximation of the global decay law, so that in the subcritical regime the global p -value will change from $1 - \theta$ at early times to $1 + \theta$ at large times. [Guo and Ogata, 1997] measured both the local and global p -values for 34 aftershocks sequences in Japan, and found that the local p -value is usually slightly larger than the global p -value [Guo and Ogata, 1997]. This is in agreement with our prediction when identifying the local p -value with $1 + \theta$ (recovered at large times) and the global p -value with $1 - \theta$ found at early times. Guo and Ogata [1997] and Ogata [1992, 1998, 2001] found a local p -value smaller than one for some aftershocks sequences in Japan. Within the confine of the ETAS model, this corresponds to the case $\theta < 0$ discussed above and in the appendix.
- The parameter K measures the rate of aftershocks triggered by each earthquake, indepen-

dently of its magnitude. Recall that the branching ratio n is proportional to K . It is usually found of the order of $K \approx 0.02$ [Ogata, 1989, 1992; Guo and Ogata, 1997], but large variations of K -value from 0.001 to 5 are reported by Ogata [1992].

- The parameter μ measures the background seismicity rate that is supposed to arise from the tectonic loading. $\mu \simeq 0$ for an aftershock sequence triggered by a single mainshock. This parameter has no influence on the branching ratio n . In real catalogs, the background seismicity only accounts for a small part of the seismic activity.

We have computed the branching ratio n and the cross-over time t^* from the ETAS parameters measured by Ogata [1989, 1992] for several seismicity sequences in Japan and elsewhere. The ETAS parameters and the n and t^* values are given in Table 1. When the b -value is not given in the text, we have computed n and t^* assuming a b -value is equal to 1. We find that the n -value is either smaller or larger than 1. This means that the seismicity can be interpreted to be either in the sub- or in the super-critical regime. An infinite n -value is found if the local p -value is smaller than one ($\theta < 0$) or if the α -value is larger than the b -value. For the same area, the ETAS parameters and the n and t^* values are found to vary in time, sometimes changing from the sub- to the super-critical regime. The characteristic time t^* shows large spatial and temporal variability, ranging from 0.4 days to 10^{22} days. Large t^* values are related to a branching ratio n close to one, *i.e.*, close to the critical point $n = 1$. The ETAS model thus provides a picture of seismicity in which sub-critical and super-critical regimes are alternating in an intermittent fashion. As we shall argue, the determination of the regime may provide important clues and quantitative tools for prediction.

Implications of the ETAS model in the sub-critical regime $n < 1$

In the sub-critical regime, the ETAS model can explain many of the departures of the global aftershock decay law from a pure Omori law.

The ETAS model contains by definition (and thus “explains”) the secondary aftershock sequences triggered by the largest aftershocks that are often observed [Correig *et al.*, 1997; Guo and Ogata, 1997; Simeonova and Solakov, 1999; Ogata, 2001]. In the

ETAS model, the fact that secondary aftershock sequences of large aftershocks can stand out above the overall background aftershock seismicity results from the factor $10^{\alpha(m_i - m_0)}$ in (1).

Our analytical results may rationalize why some alternative models of aftershock decay work better than the simple modified Omori law. In the sub-critical regime, we predict an increase of the apparent global p -value from $1 - \theta$ at early times to $1 + \theta$ at large times. To our knowledge, this change of exponent has never been observed. This change of power law may be approximated by the stretched exponential function proposed by [Kissinger, 1993; Gross and Kisslinger, 1994] to fit aftershocks sequences. In the stretched exponential model, the rate of aftershocks $\lambda(t)$ is defined by

$$\lambda(t) = K t^{q-1} e^{-(t/t_0)^q}, \quad (23)$$

where q , K and t_0 are constants. At early times, this function decays as a power law $1/t^{1-q}$ with apparent Omori’s exponent $1 - q$. For times larger than the relaxation time t_0 , the seismicity rate decays exponentially in the argument $(t/t_0)^q$. For $q < 1$, this decay is much slower than exponential and can be accounted for by an apparent power law with larger exponent. Figure 5 compares the stretched exponential function with the analytical solution of the ETAS model (16) with parameters $t^* = t_0$ and $\theta = q$, and with the Omori law of exponent $p = 1 - q$. These three laws have the same power-law behavior at early times, and then both the stretched exponential and the analytical solution (16) decay faster than the Omori law at large times. The fact that it is very difficult to distinguish the decay laws described by power laws and by stretched exponential has been illustrated in [Laherrère and Sornette, 1998] in many examples including earthquake size and fault length distributions. Kissinger [1993] and Gross and Kisslinger, 1994] compared this function to the modified Omori law $\lambda(t) = K (t + c)^{-p}$ for several aftershocks sequences in southern California. They found that the stretched exponential fit often works better for the sequences with a small p -value or a large q -value, indicative of a slow decay for small times. This is in agreement with our result that in the sub-critical regime a slowly decaying aftershock sequence (global p -value smaller than one) will then cross-over to a more rapid decay for time larger than t^* . The relaxation time t_0 ranges between 2 days and 380 days for the sequences that are better fitted by the stretched exponential [Kissinger, 1993]. This parameter is analogous to t^* found in our model, because these two

parameters define the transition from the early time power-law decay to another faster decaying behavior for large times. To further validate our results, these aftershocks sequences should be fitted using equation (16) to compare our results with the stretched exponential function and determine if the transformation of the early time power law decay is better fitted by a stretched exponential fall-off or an increase in the apparent Omori exponent from $1-\theta$ to $1+\theta$ as predicted by our results.

The ETAS model can also rationalize some correlations found empirically between seismicity parameters. It may explain the rather large variability of the global empirical p -value. *Guo and Ogata* [1995] have reported a positive correlation between the Gutenberg-Richter b -value and the p -value (exponent of the global Omori law) for several aftershocks sequences in Japan. A similar correlation has also been found by [*Kisslinger and Jones*, 1991] for several aftershock sequences in southern California, but this correlation was detectable only if the earthquake sequences were separated into thrust and strike slip events. This positive correlation between b and global p values is expected from our analysis. From equation (3), we see that a small b -value is associated with a large n value. For $n \simeq 1$, the characteristic time t^* is very large, so that the global aftershock rate decays as a power law with exponent $1-\theta$ over a large time interval. For $n > 1$ and $\theta > 0$, we see an apparent global p -value smaller than $1-\theta$ which decreases with time. In contrast, for large b -values, the branching ratio n is small and the characteristic time t^* is very small. In this case, only the large time behavior is observed with a larger exponent $1+\theta$. Consequently, in the subcritical regime, our results predict a change of the global p -value from $1-\theta$ for small b -value and times $t \ll t^*$ to $1+\theta$ for large b -values. There is also a positive correlation between p -value and b -value in the super-critical regime. For $n > 1$ or $\theta < 0$, the global aftershock sequence is characterized by an apparent exponent p smaller than $1-|\theta|$ which decreases with time. Then, we expect the apparent exponent p to be all the smaller, the smaller is the b -value, because the characteristic times t^* for $\theta > 0$ or τ for $\theta < 0$ decreases with b . The variability of the global p exponent reported by *Guo and Ogata* [1995] and *Kisslinger and Jones* [1991] may thus be explained by a change of b -value and a constant local p exponent. However, the results of *Guo and Ogata* [1997] contradict this interpretation. *Guo and Ogata* [1997] studied the same aftershocks sequences than *Guo and Ogata* [1995] but

they measured the local p -value of the ETAS model. They still found a large variability in the local p -value, and a positive correlation between this local p -value and the b -value.

Implications of the ETAS model in the super-critical regime and in the case $\theta < 0$

In the regime where the mean number of aftershocks per mainshock is larger than one (*i.e.*, $n > 1$), the mean rate of aftershocks increases exponentially for large times. However, because of the statistical fluctuations, the aftershock sequence has a finite probability to die. This probability of extinction can be evaluated for the simple branching model without time dependence [*Harris*, 1963]. Therefore, a branching ratio larger than 1 does not imply necessarily that the number of aftershocks will be infinite. If n is not too large, and if the number of aftershocks is small, there is a significant probability that the aftershock sequences will die, as observed in numerical simulations of the ETAS model. If the characteristic time t^* is very large, the aftershock sequence may not remain supercritical long enough for the exponential increase to be observed. Even if the large times exponential acceleration is rarely observed in real seismicity, it may explain the acceleration of the deformation before material failure. The early times behavior of the seismic activity preceding the exponential increase has also important possible implications for earthquake prediction, and can rationalize some empirically proposed seismic precursors, such as the low p -value [*Liu*, 1984; *Bowman*, 1997], or the relative seismic quiescence preceding large aftershocks [*Matsu'ura*, 1986; *Drakatos*, 2000].

It is widely accepted that about a third to a half of strong earthquakes are preceded by foreshocks [e.g., *Jones and Molnar*, 1979; *Bowman and Kisslinger*, 1984; *Reasenber*, 1985, 1999; *Reasenber and Jones*, 1989; *Abercombie and Mori*, 1996], *i.e.*, are preceded by an unusual high seismicity rate for time periods of the order of days to years, and distance up to hundreds kilometers. However, there is no reliable method for distinguishing foreshocks from aftershocks. Indeed, the ETAS model makes no arbitrary distinctions between foreshocks, mainshocks and aftershocks and describes all earthquakes with the same laws. While this seems a priori paradoxical, our analysis of the ETAS model provides a useful tool for identifying foreshocks, *i.e.*, earthquakes that are likely to be followed by a larger event, from usual aftershocks that are seldom followed by a larger earthquake. The

characterization of foreshocks will be performed in statistical terms rather than on a single-event basis. In other words, we will not be able to say whether any specific event is a precursor. It is the ensemble statistics that may betray a foreshock structure.

The crux of the method is that, when seismicity falls in the regime with a branching ratio $n > 1$, the corresponding earthquake sequences can be identified as foreshocks. This is because the super-critical regime corresponds to an exponentially accelerating seismicity for times larger than t^* : by a pure statistical effect, the larger number of earthquakes of any size will sample more and more the branch of the Gutenberg-Richter law toward large events. Thus by the sheer weight of numbers, larger and larger earthquakes will occur as time increases. Of course, we are not implying any precise deterministic growth law, but statistically, the largest events should indeed grow significantly, the more so, the more within the super-critical regime, the larger the branching ratio $n > 1$. Conversely, this argument implies that, in the subcritical regime, the triggered events are usual aftershocks, because a mainshock is unlikely to be followed by a larger triggered event. Foreshock sequences can thus be identified by evaluating the branching ratio n from the inversion of seismic data (times and magnitudes of an earthquake sequence) for the ETAS parameters. There is however a finite probability than a triggered event in the subcritical regime be larger than the triggering event, and thus the triggering event will be a foreshock of the triggered event. Therefore, foreshocks can be observed even in the sub-critical regime, but they are less frequent than aftershocks.

A note of caution is in order: the direct estimation of n and t^* or τ may be quite imprecise if the number of events is small. Based on our analysis and our results, the foreshock regime can be nevertheless identified with relatively good confidence if one assumes an upper bound for the local exponent p . Let us assume for instance that the local p -value is smaller than 1.3 (*i.e.*, $\theta < 0.3$); according to our results, the global exponent $p = 1 + \theta$ cannot become smaller than $1 - \theta = 0.7$ in the sub-critical regime. In contrast, in the supercritical regime, we have shown that the apparent exponent is smaller than or at most equal to $1 - \theta$. Therefore, a measure of the global p -value yielding a value smaller than 0.7, is always associated with the super-critical regime. As we said above, *Guo and Ogata* [1997] and *Ogata* [1992, 1998, 2001] found a local p -value smaller than one for some aftershocks sequences in Japan corresponding to the case $\theta < 0$. A

small global p -value can thus also result from a small local p -value. In sum, a small global p -value results either from a larger than one local p -value in the supercritical regime $n > 1$ or from a small (smaller than 1) local p -value before the exponential growth regime.

Such a small p -value precursor was first proposed empirically by *Liu* [1984], who studied several aftershocks sequences of moderate earthquakes that have been followed by a large earthquake. He proposed that a p -value smaller than 1 is a signature of a foreshock sequence, whereas $p > 1$ is associated with normal aftershock sequences with a single mainshock in the past. He suggested that p -values close to one characterize double-mainshock sequences. These empirical rules are part of the earthquake prediction method used in China [*Liu*, 1984; *Zhang et al.*, 1999]. The small precursory p -value has been used with other precursors to predict the occurrence of a $M = 6.4$ earthquake in China following an other $M = 6.4$ earthquake three months later [*Zhang et al.*, 1999]. A precursor associated with a small global p -value has also been observed by *Bowman* [1997] for a sequence in Australia. In 1987, several $M = 4 - 5$ earthquakes occurred in a region that was not seismically active before, and triggered a large number of aftershocks characterized by an abnormally low p -value of 0.3. A sequence of three $M \geq 6$ occurred one year later, followed by an aftershock sequence with a more standard p -value of 1.1. *Simeonova and Solakov* [1999] have also reported a very low p -value of 0.5, for one sequence of aftershocks in Bulgaria, that was followed one year latter by a larger earthquake. The first part of the aftershock sequence was well fitted by a modified Omori law, and then a significant deviation occurred with an abnormally high aftershock rate by comparison with the prior trend. This departure from an Omori law is expected from our results for an aftershock sequence in the super-critical regime and the very low value of the exponent p can be interpreted as the apparent exponent within the cross-over from the $1/t^{1-\theta}$ decay (14) at early times to the exponential explosion (17) at times $t > t^*$ (see Figure 3).

In addition to the small precursory p -value predicted in the regime $n > 1$, we have shown that this regime is also characterized by a decrease of the apparent global p -value with time. Such a decrease of p -value has also been identified as a precursor by *Liu* [1984].

Other patterns may be a signature of the super-critical regime. The relative precursory quiescence suggested by *Drakatos* [2000] may also be explained

by our results. In contrast to the “absolute” quiescence which detects changes in the background seismicity after removing the aftershocks from the catalog [e.g. *Wyss and Habermann, 1988*], the “relative” quiescence [*Matsu’ura, 1986; Drakatos, 2000*] takes into account the aftershocks and detects changes in seismic activity after a large mainshock by comparison with the usual Omori law decay of aftershocks. *Drakatos [2000]* studied several aftershock sequences in Greece which contains large aftershocks, *i.e.* aftershock with magnitude no smaller than $M - 1.2$, where M is the mainshock magnitude. For each sequence, he fitted the aftershock sequence by a modified Omori law up to the time of the large aftershock using a maximum likelihood method. He found that large aftershocks were often preceded by a relative quiescence by comparison with an Omori law, with an increase of the seismicity rate just before the large mainshock occurrence. Such a departure from an Omori law is predicted by our results in the super-critical regime. Indeed, in the super-critical regime, large aftershocks are likely to occur when the earthquake rate $N(t)$ changes from an Omori law to the exponential explosion for times close to t^* .

To illustrate this concept, we have performed a simulation of the ETAS model in the super-critical regime and have applied the same procedure as used by *Drakatos [2000]* to fit the synthetic aftershock sequence by an Omori law up to the time of the first large aftershock. The parameters of the synthetic catalog are $K = 0.024$, $m_0 = 0$, $c = 0.001$, $\alpha = 0.5$, $b = 0.8$ and $\theta = 0.2$, yielding $n = 1.27$ and $t^* = 4.6$. Figure 6 represents the cumulative aftershock number as a function of time for the synthetic catalog and the fit with a modified Omori law. From this figure, we see a clear relative seismic quiescence, as defined by a cumulative aftershock number smaller than that predicted by the fit. The aftershock activity recovers the level predicted by the fit at the time of the large aftershock. All these results are similar to those obtained by *Drakatos [2000]*.

In the case $n > 1$, our results predict an exponential increase of the seismicity rate at large times. Because we assume that the magnitude distribution is independent of time, the same exponential acceleration is expected for both the cumulative energy release and the cumulative number of earthquakes. *Sykes and Jaumé [1990]* found that several large earthquakes in the San-Francisco Bay area were preceded by an acceleration of the cumulative energy release that can be fitted by an exponential function, as

predicted by our results. In laboratory experiments of rupture, several studies have also observed an exponential acceleration of the seismic energy release before the macroscopic rupture [*Scholz, 1968; Meredith et al., 1990; Main et al., 1992*].

More recently, many studies have reported an acceleration of seismicity prior to great events (see [*Sammis and Sornette, 2001; Vere-Jones et al., 2001*] for reviews) but they used a power-law instead of an exponential law to fit the acceleration of seismicity. A power-law increase of the seismicity before rupture is predicted by several statistical models of rupture in heterogeneous media, which consider the global rupture or the great earthquake as a critical point (see [*Sornette, 2000a*] for a review). Note that it is often difficult to distinguish in real data an exponential increase from a power-law increase, especially with a small number of points and for times far from the rupture time. No systematic study has been undertaken that compares these two laws to test if the acceleration of the seismicity is better fitted by a power-law rather than by an exponential law (see however [*Johansen et al., 1996*]).

We have stressed that the ETAS model is fundamentally a mean field approximation (branching process) which neglects “loops”, *i.e.*, multiple interactions (see Figure 1). An important consequence of this approximation is that the super-critical regime cannot lead to a growth rate faster than exponential. Indeed, recall that an exponential growth is characterized by a time derivative of the number of events proportional to the number of events $dN/dt = N/t^*$, *i.e.*, is fundamentally a *linear* process. In a sequel to the present work [*Sornette and Helmstetter, 2001*], we show however that for $b < \alpha$, the impact of the largest earthquake induces an effective nonlinearity which leads to a faster-than-exponential growth rate, possibly leading to a finite-time singularity [*Sammis and Sornette, 2002*]. A faster-than-exponential growth rate may also be obtained by introducing multiple interactions between earthquakes and positive feedback: rather than the linear law $dN/dt = N/t^*$ expressing the condition that each “daughter” has only one “mother”, we may expect an effective law $dN/dt \sim N^\delta$, with $\delta > 1$ providing a measure of the effective number of ancestors impacting directly on the birth of a daughter. We may thus expect that an improvement of the ETAS model beyond the “mean-field” approximation would lead to power law acceleration of seismicity.

Other precursory patterns may also be related to

the super-critical regime: they comprise the precursory earthquake swarm or burst of aftershocks [Evison, 1977; Keilis-Borok *et al.*, 1980a, 1980b; Molchan *et al.*, 1990; Evison and Rhoades, 1999]. Swarms are earthquake sequences characterized by high clustering in space and time and the occurrence of several large events with magnitude larger than $M - 1$, where M is the magnitude of the largest event. A burst of aftershocks is a sequence of one or more mainshocks with abnormally large number of aftershocks at the beginning of their aftershock sequences [Keilis-Borok *et al.*, 1980a]. From our results, an abnormally high aftershock rate or a sequence with several large events are expected in the super-critical regime.

Temporal change of n -value and transition from one regime to the other one

It is often reported that the b and p values vary in space and time [e.g., Smith, 1981; Guo and Ogata, 1995, 1997; Wiemer and Katsumata, 1999]. We have documented that a part of the observed variation of the exponent p may not be genuine but result from an inadequate parameterization of a more complex reality. Because n and t^* are function of b , p and the other ETAS parameters, we expect the fundamental parameters of the ETAS model, namely n and t^* , to vary significantly in space and time. The branching ratio n plays the role of a “control” parameter quantifying the distance from the critical point $n = 1$ between the sub-critical and the super-critical regime; t^* is a crossover time and is sensitive to details of the systems. As a consequence, it is very reasonable to expect that the Earth’s crust will change from the sub-critical to the super-critical regime and vice-versa, as a function of time and location.

Equation (3) shows that the branching ratio n is a decreasing function of b . Accordingly, this may rationalize the observation that large earthquakes are sometimes preceded by a decrease of the b -value [e.g. Smith, 1981]. A decrease of the b -value leads to an increase of the n -value, that can move the seismicity from the sub-critical to the super-critical regime, and thus increase the probability to observe a large earthquake. Other ETAS parameters (α , K , p and c) may also change in time and move the seismicity from one regime to the other one. Ogata [1989] measured the ETAS model parameters before and after the 1984 Western Nagano Prefecture earthquake ($M = 6.8$). He found that the seismic activity preceding the mainshock was characterized by a lower b , c , K parameters and local p values than the seismicity

following the mainshock. He also obtained a larger α -value for the seismicity preceding the mainshock. All these changes of parameters, except the change in K , lead to a larger n -value before the mainshock than after. Before the mainshock, n is in principle infinite because the local p -value is smaller than one. As we already discussed, this corresponds to an explosive super-critical regime of growing seismicity. After the mainshock, we find $n = 0.92$ and $t^* = 10^6$ days, using the determination of the ETAS parameters. The seismicity has thus changed from a super-critical regime before the mainshock to a sub-critical regime after the mainshock.

Conclusion

We have provided analytical solutions of the ETAS model, which describes foreshocks, aftershocks and mainshocks on the same footing. Each event triggers an aftershock sequence with a rate that decays according to the local Omori law with an exponent $p = 1 + \theta$. The number of aftershocks per event increases with its magnitude. We suggest that the Earth’s crust at a given time and location may be characterized by its branching ratio n , quantifying its regime. We propose that n is a fundamental parameter for understanding and characterizing the organization of the seismicity within the Earth’s crust. In the sub-critical regime ($n < 1$), the global rate of aftershocks (including secondary aftershocks) decays with the time from the mainshock with a decay law different from the local Omori law. We find a crossover from an Omori exponent $1 - \theta$ for $t < t^*$ to $1 + \theta$ for $t > t^*$. The modified Omori law is thus only an approximation of the decay law of the global aftershock sequence. In the super-critical regime ($n > 1$ and $\theta > 0$), we find a novel transition from an Omori decay law with an exponent $1 - \theta$ at early times to an explosive exponential increase of the seismicity rate at large times. The case $\theta < 0$ leads to an infinite n -value, due to the slow decay with time of the local Omori law. In this case, we find a transition from an Omori law with exponent $1 - |\theta|$ similar to the local law, to an exponential increase at large times, with a crossover time τ different from the characteristic time t^* found in the case $\theta > 0$. These results can rationalize many of the stylized facts reported for foreshock and aftershock sequences, such as the suggestion that a small p -value may be a precursor of a large earthquake, the relative seismic quiescence preceding large aftershocks, the positive correlation between b and p -values, the observation that great earthquakes are

sometimes preceded by a decrease of b -value and the acceleration of the seismicity preceding great earthquakes.

Finally, we would like to mention that our analysis can be generalized to various other choices of the local Omori law and of the magnitude distribution. The ETAS model can also be extended to describe the spatial distribution of the seismicity [Helmstetter and Sornette, 2002].

Acknowledgments. We are very grateful to J.R. Grasso, G. Ouillon, V. Pisarenko, A. Sornette and D. Vere-Jones for useful suggestions and discussions and to D. Vere-Jones as a referee for his constructive remarks and for pointing out the relevant mathematical literature on “self-exciting” point processes. We thank Y. Ogata for kindly providing a copy of Ramselaar’s Master thesis.

Appendix: technical derivation of the analytical solution

In this appendix, we provide the technical derivation of the results used in the main text for the sub-critical and super-critical regimes. We start from equation (9).

General derivation for $\theta > 0$

The integral over τ is the convolution of $\lambda_{m'}$ with $\phi_{m'}$. Since there is an origin of time and we have a convolution operator, the natural tool is the Laplace transform $\hat{f}(\beta) \equiv \int_0^{+\infty} f(t)e^{-\beta t} dt$. Applying the Laplace transform to (9) yields

$$\hat{\lambda}_m(\beta) = \hat{S}(\beta, m) + P(m) \int_{m_0}^{\infty} dm' \hat{\phi}_{m'}(\beta) \hat{\lambda}_{m'}(\beta). \quad (\text{A1})$$

where the r.h.s. has used the convolution theorem that the Laplace transform of a convolution of two functions is the product of the Laplace transform of the two functions. Let us now apply the integral operator $\int_{m_0}^{\infty} dm \hat{\phi}_m(\beta)$ on both sides of (A1) and define

$$\lambda(\beta) \equiv \int_{m_0}^{\infty} dm \hat{\phi}_m(\beta) \hat{\lambda}_m(\beta), \quad (\text{A2})$$

$$Q(\beta) \equiv \int_{m_0}^{\infty} dm \hat{\phi}_m(\beta) P(m), \quad (\text{A3})$$

and

$$\mathcal{S}(\beta) \equiv \int_{m_0}^{\infty} dm \hat{\phi}_m(\beta) \hat{S}(\beta, m). \quad (\text{A4})$$

Then, expression (A1) yields

$$\lambda(\beta) = \mathcal{S}(\beta) + Q(\beta)\lambda(\beta), \quad (\text{A5})$$

whose solution is

$$\lambda(\beta) = \frac{\mathcal{S}(\beta)}{1 - Q(\beta)}. \quad (\text{A6})$$

This expression gives $\lambda_m(t)$ after inversion of the integral operator $\int_{m_0}^{\infty} dm \hat{\phi}_m(\beta)$ and of the Laplace transform.

The key quantity controlling the dependence of $\lambda_m(t)$ is

$$Q(\beta) = \frac{K}{\theta c^\theta} \left(\int_{m_0}^{\infty} dm 10^{\alpha(m-m_0)} P(m) \right) \left(\theta \int_0^{\infty} dt \frac{e^{-\beta ct}}{(t+1)^{1+\theta}} \right), \quad (\text{A7})$$

obtained by replacing the expression of $\phi_m(t)$ defined in (1) and normalizing $t/c \rightarrow t$. Using $P(m) = \ln(10) b 10^{-b(m-m_0)}$, we obtain

$$Q(\beta) = n R(\beta c), \quad (\text{A8})$$

where we have used the expression (3) of n and defined

$$R(\beta) \equiv \theta \int_0^{\infty} dt \frac{e^{-\beta t}}{(t+1)^{1+\theta}} = \theta e^\beta \beta^\theta \Gamma(-\theta, \beta) = 1 - e^\beta \beta^\theta \Gamma(1-\theta, \beta), \quad (\text{A9})$$

where

$$\Gamma(a, x) = \int_x^{\infty} dt e^{-t} t^{a-1} \quad (\text{A10})$$

is the (complementary) incomplete Gamma function [Abramowitz and Stegun, 1964] and we have used $\Gamma(1+a, x) = a\Gamma(a, x) + x^a e^{-x}$ obtained by integration by part. Using the expansion of the incomplete Gamma function [Olver, 1974]

$$\Gamma(a, x) = \Gamma(a) - \sum_{k=0}^{+\infty} \frac{(-1)^k x^{a+k}}{k! (a+k)}, \quad \text{for } a > 0, \quad (\text{A11})$$

we obtain

$$R(\beta) = 1 - \Gamma(1-\theta) \beta^\theta + \frac{1}{1-\theta} \beta + \mathcal{O}(\beta^{1+\theta}, \beta^2, \beta^{2+\theta}, \beta^3, \dots). \quad (\text{A12})$$

It is possible, using the full expansion of the incomplete Gamma function, to estimate the value of $\lambda(\beta)$ when the second term $\frac{1}{1-\theta} \beta$ of the expansion cannot be neglected anymore compared with the term proportional to β^θ . Thus, the expansion (A12) using the first two terms only $R(\beta) = 1 - \Gamma(1-\theta) \beta^\theta$ becomes invalid for $\beta > [\Gamma(1-\theta)(1-\theta)]^{1/(1-\theta)}$, i.e., for times smaller than $[\Gamma(2-\theta)]^{-1/(1-\theta)}$. For all practical purpose, this is a small value and we can use safely the expansion (A12) in the following calculations.

Let us now make explicit $\lambda(\beta)$:

$$\lambda(\beta) = \frac{K}{\theta c^\theta} R(\beta c) \int_{m_0}^{\infty} dm 10^{\alpha(m-m_0)} \int_0^{\infty} dt \lambda_m(t) e^{-\beta t}. \quad (\text{A13})$$

Using the definition of $\lambda(t)$ given by (11) and the factorization of the times and magnitudes in (A13), we obtain

$$\lambda(\beta) = n R(\beta c) \hat{\lambda}(\beta), \quad (\text{A14})$$

where

$$\hat{\lambda}(\beta) = \int_0^{\infty} dt \lambda(t) e^{-\beta t}. \quad (\text{A15})$$

Replacing (A14) in (A6) gives

$$\hat{\lambda}(\beta) = \frac{\mathcal{S}(\beta)}{nR(\beta c)(1 - nR(\beta c))}. \quad (\text{A16})$$

When a great earthquake occurs at the origin of time $t = 0$ with magnitude M , $S(t, m) = \delta(t) \delta(m - M)$, expression (A4) gives

$$\mathcal{S}(\beta) = \frac{K}{\theta c^\theta} 10^{\alpha(M-m_0)} R(\beta c). \quad (\text{A17})$$

Thus, expression (A16) becomes

$$\hat{\lambda}(\beta) = \frac{b - \alpha}{b} \frac{10^{\alpha(M-m_0)}}{(1 - nR(\beta c))}. \quad (\text{A18})$$

The dependence of $\hat{\lambda}(\beta)$ on β is uniquely controlled by the denominator $1 - nR(\beta c)$.

The sub-critical regime $n < 1$

The analysis proceeds exactly as in [Sornette and Sornette, 1999]. For $0 < \theta < 1$, and for small β (large times), $\hat{\lambda}(\beta)$ given by (A18) is

$$\hat{\lambda}(\beta) = \frac{S_0}{1 - n[1 - d(\beta c)^\theta]} = \frac{S_0}{(1 - n)} \left(\frac{1}{1 + (\beta t^*)^\theta} \right), \quad (\text{A19})$$

where t^* is defined by (12) and the external source term S_0 is defined by (13). We retrieve equation (13) of [Sornette and Sornette, 1999] with the correspondence $t_0 \rightarrow c$.

Two cases must be distinguished.

- $\beta t^* < 1$ corresponds to $t > t^*$ by identifying as usual the dual variable β to t in the Laplace transform with $1/t$. In this case, we can expand $\frac{1}{1 + (\beta t^*)^\theta}$, which leads to

$$\hat{\lambda}_{t > t^*}(\beta) \sim \frac{S_0}{1 - n} [1 - (\beta t^*)^\theta]. \quad (\text{A20})$$

We recognize the Laplace transform of a power law of exponent θ , *i.e.*

$$\lambda_{t > t^*}(t) \sim \frac{S_0}{\Gamma(\theta)(1 - n)} \frac{t^{*\theta}}{t^{1+\theta}} \quad \text{for } t > t^*. \quad (\text{A21})$$

- For $t < t^*$, $\beta t^* > 1$ and (A19) can be written with a good approximation as

$$\hat{\lambda}_{t < t^*}(\beta) = \frac{S_0}{(1 - n)(\beta t^*)^\theta} \sim \beta^{-\theta}. \quad (\text{A22})$$

Denoting $\Gamma(z) \equiv \int_0^{+\infty} dt e^{-t} t^{z-1}$, we see that $\int_0^{+\infty} dt e^{-\beta t} t^{z-1} = \Gamma(z)\beta^{-z}$. Comparing with (A22), we thus get

$$\lambda_{t < t^*}(t) \sim \frac{S_0}{\Gamma(\theta)(1 - n)} \frac{t^{*\theta}}{t^{1-\theta}} \quad \text{for } t < t^*. \quad (\text{A23})$$

We verify the self-consistency of the two solutions $\lambda_{t > t^*}(t)$ and $\lambda_{t < t^*}(t)$ by checking that $\lambda_{t > t^*}(t^*) = \lambda_{t < t^*}(t^*)$.

In other words, t^* is indeed the transition time at which the “short-time” regime $\lambda_{t < t^*}(t)$ crosses over to the “long-time” regime $\lambda_{t > t^*}(t)$.

We now calculate the full expression of $\lambda(t)$ valid at all times. We expand

$$\frac{1}{(\beta t^*)^\theta + 1} = \frac{1}{(\beta t^*)^\theta} \frac{1}{(\beta t^*)^{-\theta} + 1} = \frac{1}{(\beta t^*)^\theta} \sum_{k=0}^{\infty} (-1)^k (\beta t^*)^{-k\theta}, \quad (\text{A24})$$

Thus, by taking the inverse Laplace transform

$$\lambda(t) = \frac{S_0}{1 - n} \frac{1}{2\pi i} \int_{c-i\infty}^{c+i\infty} d\beta e^{\beta t} \sum_{k=0}^{\infty} (-1)^k (\beta t^*)^{-(k+1)\theta}. \quad (\text{A25})$$

The inverse Laplace transform of $\beta^{-(k+1)\theta}$ is $t^{(k+1)\theta-1}/\Gamma((k+1)\theta)$. This allows us to write

$$\lambda(t) = \frac{S_0}{1 - n} \frac{t^{*\theta}}{t^{1-\theta}} \sum_{k=0}^{\infty} (-1)^k \frac{(t/t^*)^{k\theta}}{\Gamma((k+1)\theta)} \quad (\text{A26})$$

Expression (A26) provides the solution that describes the cross-over from the $1/t^{1-\theta}$ Omori’s law (A23) at early times to the $1/t^{1+\theta}$ Omori’s law (A21) at large times. The series $\sum_{k=0}^{\infty} (-1)^k \frac{(t/t^*)^{k\theta}}{\Gamma((k+1)\theta)}$ is a series representation of the special Fox function [Glöckle and Nonnenmacher, 1993] and it is also related to the generalized Mittag-Leffler function.

For large times $t \gg t^*$, a direct numerical evaluation of $\lambda(t)$ from equation (A26) is impossible due to the very slow convergence of the series. The padé summation method [Bender and Orszag, 1978] can be used to improve the convergence of this series and to evaluate numerically (A26) for all times.

The super-critical regime $n > 1$

We can analyze this regime by putting $n > 1$ in (A18) which can be written under a form similar to (A19):

$$\hat{\lambda}(\beta) = \frac{S_0}{(1 - nR(\beta c))} = \frac{S_0}{dn(\beta c)^\theta - (n - 1)} = \frac{S_0}{(n - 1)} \left(\frac{1}{(\beta t^*)^\theta - 1} \right), \quad (\text{A27})$$

In the second and third equalities of (A27), we have used the small β -expansion (A12) of $R(\beta c)$ valid for $0 < \theta < 1$.

At early times $c \ll t \ll t^*$, *i.e.*, $\beta t^* \gg 1$, $\hat{\lambda}(\beta) \approx \frac{S_0}{(n-1)(\beta t^*)^\theta}$ which is the Laplace transform of (A23): thus, the early decay rate of aftershocks is the same $\sim 1/t^{1-\theta}$ as for the sub-critical regime (A23). However, as time increases, the dual β of t decreases and $\hat{\lambda}(\beta)$ grows faster than $\sim (\beta c)^{-\theta}$ due to the presence of the negative term $-(n - 1)$. This can be seen as an apparent exponent $\theta_{\text{app}} > \theta$ increasing progressively such that $dn(\beta t^*)^\theta - 1 \approx C(\beta t^*)^{\theta_{\text{app}}}$, where C is a constant. Note that $\theta_{\text{app}} > \theta$ for the pure power law $C(\beta c)^{\theta_{\text{app}}}$ to mimic the acceleration induced by the negative correction $-(n - 1)$. The seismic rate will thus decay approximately as $\sim 1/t^{1-\theta_{\text{app}}(t)}$.

The large time behavior is controlled by the pole at $\beta = 1/t^*$ of $\hat{\lambda}(\beta)$. Close to $1/t^*$,

$$\hat{\lambda}(\beta) \approx \frac{S_0}{(n-1)\theta} \frac{1}{\beta t^* - 1}. \quad (\text{A28})$$

The inverse Laplace transform is thus

$$\lambda(t) = (2\pi i)^{-1} \int_{c-i\infty}^{c+i\infty} d\beta e^{\beta t} \hat{\lambda}(\beta) \sim \frac{S_0}{(n-1)t^*\theta} e^{t/t^*} \quad (\text{A29})$$

exhibiting the exponential growth at large times. Expression (12) shows that $1/t^* \sim |1-n|^{1/\theta}$. Thus, as expected, the exponential growth disappears as $n \rightarrow 1^+$.

We now calculate the full expression of $\lambda(t)$ valid at all times. We expand

$$\frac{1}{(\beta t^*)^\theta - 1} = \frac{1}{(\beta t^*)^\theta} \frac{1}{1 - (\beta t^*)^{-\theta}} = \frac{1}{(\beta t^*)^\theta} \sum_{k=0}^{\infty} (\beta t^*)^{-k\theta}, \quad (\text{A30})$$

Thus

$$\lambda(t) = \frac{S_0}{(n-1)} \frac{1}{2\pi i} \int_{c-i\infty}^{c+i\infty} d\beta e^{\beta t} \sum_{k=0}^{\infty} (\beta t^*)^{-(k+1)\theta}. \quad (\text{A31})$$

The inverse Laplace transform of $1/\beta^{(k+1)\theta}$ is $t^{(k+1)\theta-1}/\Gamma((k+1)\theta)$. This allows us to write

$$\lambda(t) = \frac{S_0}{(n-1)} \frac{t^{*-\theta}}{t^{1-\theta}} \sum_{k=0}^{\infty} \frac{(t/t^*)^{k\theta}}{\Gamma((k+1)\theta)} \quad (\text{A32})$$

Expression (A32) provides the solution that describes the cross-over from the $1/t^{1-\theta}$ Omori's law at early times to the exponential growth at large times. Note that the solution (A32) can be obtained directly from (A26) by removing the alternating sign $(-1)^k$ in the sum. The solution (A32) retrieves the two regimes discussed before.

1. For $t < t^*$, the sum in (A32) is close to $1/\Gamma(\theta)$, which leads to

$$\lambda(t) \approx \frac{S_0}{\Gamma(\theta)(n-1)} \frac{t^{*-\theta}}{t^{1-\theta}}. \quad (\text{A33})$$

2. For $t \geq t^*$, the sum dominates. The sum is very similar to the series expansion of e^{t/t^*} and is actually proportional to e^{t/t^*} for large t . This result is obvious for $\theta = 1$ since the series expansion becomes identical to that of e^{t/t^*} . This can be justified for other values of θ as follows. For $\theta \rightarrow 0$, the discrete sum transforms into a continuous integral of the type

$$\int_0^{\infty} dx t^x / \Gamma(x). \quad (\text{A34})$$

A saddle-node approximation, performed using the Stirling approximation (which already gives a very

good precision for small z) $\Gamma(z) \approx \sqrt{2\pi} e^{-z} z^{z-\frac{1}{2}}$, shows that the saddle-node of the integrand occurs for $x \approx t/t^*$, which then gives $\lambda(t) \sim e^{t/t^*}$. For arbitrary θ , we can use the Poisson's summation rule

$$\sum_{r=-\infty}^{+\infty} f(r) = \int_{-\infty}^{+\infty} du f(u) + \sum_{q=1}^{+\infty} \int_{-\infty}^{+\infty} du f(u) \cos[2\pi qu], \quad (\text{A35})$$

on the function defined by

$$f(r) \equiv \frac{(t/t^*)^{r\theta}}{\Gamma(r\theta + \theta)}, \quad \text{for } r \geq 0 \quad (\text{A36})$$

and $f(r) = 0$ for $r < 0$. The left-hand-side of (A35) is nothing but the semi-infinite sum in (A32). The first term in the right-hand-side retrieves the integral (A34) encountered for the case $\theta \rightarrow 0$. This term thus contributes a term proportional to e^{t/t^*} . All the other terms contribute negative powers of t and are thus negligible compared to the exponential for $t > t^*$. This can be seen from the fact that each term with $q \geq 1$ is similar to the sum in (A26) for the subcritical case with alternating signs. The larger q is, the faster is the frequency of the leading dependence $\lambda(t) \sim e^{t/t^*}$ valid for any $0 \leq \theta \leq 1$ retrieves the limiting behavior already given in (A29) from a different approach for large times $t \gg t^*$. It has also been proved rigorously in [Ramselaar, 1990].

Case $\theta < 0$ corresponding to a local Omori's law exponent $p < 1$

The general equation (9) still holds in this case and the general derivation starting with (A1) up to (A6) still applies. The key quantity controlling the dependence of $\lambda_m(t)$ is still $Q(\beta)$ defined by (A7). Writing $\theta = -|\theta|$, we have

$$Q(\beta) = n_0 R'(\beta c), \quad (\text{A37})$$

where n_0 is defined by (4) and

$$R'(\beta) \equiv \int_0^{\infty} dt \frac{e^{-\beta t}}{(t+1)^{1-|\theta|}} = e^\beta \beta^{-|\theta|} \Gamma(|\theta|, \beta) \quad (\text{A38})$$

where $\Gamma(a, x)$ is the (complementary) incomplete Gamma function defined by (A10). Using the exact expansion (A11), we obtain

$$Q(\beta) = n_0 e^{\beta c} (\beta c)^{-|\theta|} \left(\Gamma(|\theta|) - \sum_{k=0}^{+\infty} \frac{(-1)^k (\beta c)^{|\theta|+k}}{k! (|\theta|+k)} \right). \quad (\text{A39})$$

For small β 's (*i.e.*, large times), expression (A39) has the following leading behavior

$$Q(\beta) = n_0 \Gamma(|\theta|) (\beta c)^{-|\theta|} - \frac{n_0}{|\theta|} + n_0 \Gamma(|\theta|) (\beta c)^{1-|\theta|} + h.o.t. \quad (\text{A40})$$

where *h.o.t.* stands for higher-order terms in the expansion in increasing powers of βc .

The source term $\mathcal{S}(\beta)$ in the denominator of $\hat{\lambda}(\beta)$ given by (A6) is now given by

$$\mathcal{S}(\beta) = K c^{|\theta|} 10^{\alpha(M-m_0)} R'(\beta c) . \quad (\text{A41})$$

Expression (A6) for $\hat{\lambda}(\beta)$ then yields

$$\hat{\lambda}(\beta) = \frac{S_0}{1 - Q(\beta c)} , \quad (\text{A42})$$

where $R'(\beta c)$ is given by (A38), n_0 is defined by (4) and S_0 is defined by (13). The dependence of $\hat{\lambda}(\beta)$ on β is uniquely controlled by the denominator $1 - Q(\beta c) = 1 - n_0 R'(\beta c)$.

Using (A40), we get the leading behavior for small βc

$$\begin{aligned} \hat{\lambda}(\beta) &= \frac{S_0}{1 + \frac{n_0}{|\theta|} - n_0 \Gamma(|\theta|) (\beta c)^{-|\theta|}} \\ &= \frac{S_0}{\left(1 + \frac{n_0}{|\theta|}\right) (1 - (\beta \tau)^{-|\theta|})} \end{aligned} \quad (\text{A43})$$

where the characteristic time τ is given by (19).

At early times $c < t < \tau$, $(\beta \tau)^{-|\theta|} < 1$ so that

$$\hat{\lambda}(\beta) \approx \frac{S_0}{\left(1 + \frac{n_0}{|\theta|}\right)} (1 + (\tau \beta)^{-|\theta|}) . \quad (\text{A44})$$

By applying the inverse Laplace transform, the constant term contributes a Dirac function $\delta(t)$ which is irrelevant as the calculation is valid only for $t > c$. The other term $(\tau \beta)^{-|\theta|}$ gives

$$\lambda(t) = \frac{S_0}{\left(1 + \frac{n_0}{|\theta|}\right) \Gamma(|\theta|)} \frac{\tau^{-|\theta|}}{t^{1-|\theta|}} . \quad (\text{A45})$$

The early time behavior of $\lambda(t)$ is thus similar to the local Omori law $1/t^{1-|\theta|}$.

Similarly to the super-critical case $n > 1$ of the regime $\theta > 0$, the long time dependence of the regime $\theta < 0$ is controlled by a simple pole $\beta^* = \frac{1}{\tau}$.

Thus, the long-time seismicity is given by

$$\lambda(t) = \frac{S_0}{\left(1 + \frac{n_0}{|\theta|}\right) \tau^{|\theta|}} e^{t/\tau} \quad (\text{A46})$$

We can also calculate the full expression of $\lambda(t)$ valid at all times $t > c$. We expand

$$\frac{1}{1 - (\beta \tau)^{-|\theta|}} = \sum_{k=0}^{\infty} (\beta \tau)^{-k|\theta|} , \quad (\text{A47})$$

Removing the constant term, which by the inverse Laplace transform contributes a Dirac function $\delta(t)$ which is irrelevant as the calculation is valid only for $t \gg c$, we get

$$\lambda(t) = \frac{S_0}{\left(1 + \frac{n_0}{|\theta|}\right)} \frac{1}{2\pi i} \int_{c-i\infty}^{c+i\infty} d\beta e^{\beta t} \sum_{k=1}^{\infty} (\beta \tau)^{-k|\theta|} . \quad (\text{A48})$$

The inverse Laplace transform of $1/\beta^{k|\theta|}$ is $t^{k|\theta|-1}/\Gamma(k|\theta|)$. This allows us to write

$$\lambda(t) = \frac{S_0}{\left(1 + \frac{n_0}{|\theta|}\right)} \frac{1}{t} \sum_{k=1}^{\infty} \frac{(t/\tau)^{k|\theta|}}{\Gamma(k|\theta|)} \quad (\text{A49})$$

Expression (A49) provides the solution that describes the cross-over from the local Omori law $1/t^{1-|\theta|}$ at early times to the exponential growth at large times.

References

- Abercrombie, R. E. and J. Mori, Occurrence patterns of foreshocks to large earthquakes in the western United States, *Nature*, 381, 1996.
- Abramowitz, M. and I. A. Stegun, *Handbook of Mathematical Functions* Dover Publications, New York, 1964.
- Bender, C. M. and S. Orzag, *Advanced Mathematical Methods for Scientists and Engineers*, Mc. Graw-Hill, 1978.
- Blanter, E. M., M.G. Shnirman, J.-L. Le Mouél and C. J. Allègre, Scaling laws in blocks dynamics and dynamic self-organized criticality, *Phys. Earth Planet. Int.*, 99, 295-307, 1997.
- Bowman, J. R., A seismicity precursor to a sequence of Ms 6.3-6.7 midplate earthquakes in Australia, *Pure Appl. Geophys.*, 149, 61-78, 1997.
- Bowman, J. R. and K. Kisslinger, A test of foreshocks occurrence in the central Aleutian island arc, *Bull. Seism. Soc. Am.*, 74, 181-197, 1984.
- Console, R. and M. Murru, A simple and testable model for earthquake clustering, *J. Geophys. Res.*, 166, 8699-8711, 2001.
- Correig, A. M., M. Urquizú and J. Vila, Aftershock series of event February 18, 1996: An interpretation in terms of self-organized criticality, *J. Geophys. Res.*, 102, 27,407-27,420, 1997.
- Daley, D. J. and D. Vere-Jones, *An Introduction to the Theory of Point Processes*, Springer, 1988.
- Davis, S. D. and C. Frohlich, Single-link cluster analysis of earthquake aftershocks: decay laws and regional variations, *J. Geophys. Res.*, 96, 6335-6350, 1991.
- Drakatos, G., Relative seismic quiescence before large aftershocks, *Pure Appl. Geophys.*, 157, 1407-1421, 2000.
- Dreger, D. and B. Savage, Aftershocks of the 1952 Kern County, California, earthquake sequence, *Bull. Seism. Soc. Am.*, 89, 1094-1108, 1999.
- Evison, F. F., The precursory earthquake swarm, *Phys. Earth Planet. Inter.*, 15, 19-23, 1977.
- Evison, F. F. and D. A. Rhoades, The precursory earthquake swarm in Japan: hypothesis test, *Earth Planets and Space*, 51, 1267-1277, 1999.
- Feller, W. *An Introduction to Probability Theory and its Applications*, vol II, John Wiley and sons, New-York, 1971.
- Felzer, K. R., T. W. Becker, R. E. Abercrombie, G. Ekström and J.R. Rice, Triggering of the 1999 M_W 7.1

- Hector Mine earthquake by aftershocks of the 1992 M_W 7.3 Landers earthquake, submitted to *J. Geophys. Res.*, 2001.
- Gardner, J. K. and L. Knopoff, Is the sequence of earthquakes in Southern California, with aftershocks removed, Poissonian?, *Bull. Seismol. Soc. Amer.*, *64*, 1363-1367, 1974.
- Glöckle, W. and T. F. Nonnenmacher, Fox function representation of non-Debye relaxation processes, *J. Stat. Phys.*, *71*, 1993.
- Gross, S. and C. Kisslinger, Tests of models of aftershocks rate decay, *Bull. Seism. Soc. Am.*, *84*, 1571-1579, 1994.
- Guo, Z. and Y. Ogata, Correlations between characteristic parameters of aftershock distributions in time, space and magnitude, *Geophys. Res. Lett.*, *22*, 993-996, 1995.
- Guo, Z. and Y. Ogata, Statistical relations between the parameters of aftershocks in time, space and magnitude, *J. Geophys. Res.*, *102*, 2857-2873, 1997.
- Harris, T. E., *The theory of branching processes*, Springer, Berlin, 1963.
- Harris, R. A., *Stress triggers, stress shadows, and seismic hazard*, *International Handbook of Earthquake and Engineering Seismology*, W.H.K. Lee, H. Kanamori, P. C. Jennings, C. Kisslinger, Eds., Chapter 73, 48 pages, 2001.
- Hawkes, A. G., Point spectra of some mutually exciting point processes. *Journal of Royal Statistical Society, series B*, *33*, 438-443, 1971.
- Hawkes, A. G., Spectra of some mutually exciting point processes with associated variables, In *Stochastic Point Processes*, ed. P.A.W. Lewis, Wiley, 261-271, 1972.
- Hawkes, A. G. and Adamopoulos, L., Cluster models for earthquakes - regional comparisons, *Bull. Internat. Stat. Inst.*, *45*, 454-461, 1973.
- Hawkes, A. G. and Oakes D., A cluster representation of a self-exciting process, *Journal Appl. Prob.*, *11*, 493-503, 1974.
- Helmstetter, A. and D. Sornette, Diffusion of aftershocks epicenters and Omori's law: exact mapping to generalized continuous time random walk models, submitted to *Phys. Rev. E*, 2002 (e-print at <http://xxx.lanl.gov/abs/cond-mat/0203505>)
- Hough, S. E. and L. M. Jones, Aftershocks; Are they earthquakes or afterthoughts ? *EOS, Transactions, American Geophysical Union*, *78*, N45, 505-508, 1997.
- Huang, Y., H. Saleur, C. G. Sammis, D. Sornette, Precursors, aftershocks, criticality and self-organized criticality, *Europhysics Letters*, *41*, 43-48, 1998.
- Johansen, A., D. Sornette, H. Wakita, U. Tsunogai, W. I. Newman and H. Saleur, Discrete scaling in earthquake precursor phenomena : evidence in the Kobe earthquake, Japan, *J.Phys.I France*, *6*, 1391-1402, 1996.
- Jones, L. M. and P. Molnar, Some characteristics of foreshocks and their possible relationship to earthquake prediction and premonitory slip on faults, *J. Geophys. Res.*, *84*, 3596-3608, 1979.
- Kagan, Y. Y., Likelihood analysis of earthquake catalogues *Geophys. J. Int.*, *106*, 135-148, 1991
- Kagan, Y. Y. and D. D. Jackson, Spatial aftershock distribution: Effect of normal stress, *J. Geophys. Res.*, *103*, 24453-24467, 1998.
- Kagan, Y. Y. and D. D. Jackson, Probabilistic forecasting of earthquakes, *Geophys. J. Int.*, *143*, 438-453, 2000.
- Kagan, Y. Y. and L. Knopoff, Stochastic synthesis of earthquake catalogs, *J. Geophys. Res.*, *86*, 2853-2862, 1981.
- Kagan, Y. Y. and L. Knopoff, Statistical short-term earthquake prediction, *Science*, *236*, 1563-1467, 1987.
- Keilis-Borok, V. I., L. Knopoff, I.M. Rotvain and T.M. Sidorenko, Burst of seismicity as long-term precursors of strong earthquake, *J. Geophys. Res.*, *85*, 803-811, 1980a.
- Keilis-Borok, V. I., L. Knopoff and C.R. Allen, Long-term premonitory seismicity patterns in Tibet and the Himalayas, *J. Geophys. Res.*, *85*, 813-820, 1980b.
- Kisslinger, C., The stretched exponential function as an alternative model for aftershock decay rate, *J. Geophys. Res.*, *98*, 1913-1921, 1993.
- Kisslinger, C. and L. M. Jones, Properties of aftershocks sequences in Southern California, *J. Geophys. Res.*, *96*, 11947-11958, 1991.
- Knopoff, L., Y. Y. Kagan and R. Knopoff, b -values for fore- and aftershocks in real and simulated earthquakes sequences, *Bull. Seism. Soc. Am.*, *72*, 1663-1676, 1982.
- Laherrère, J. and D. Sornette, Stretched exponential distributions in Nature and Economy: "Fat tails" with characteristic scales, *European Physical Journal B*, *2*, 525-539, 1998.
- Lee, M. W. and D. Sornette, Novel mechanism for discrete scale invariance in sandpile models, *European Physical Journal B*, *15*, 193-197, 2000.
- Liu, Z. R., Earthquake frequency and prediction, *Bull. Seism. Soc. Am.*, *74*, 255-265, 1984.
- Main, I. G., P. G. Meredith and P. R. Sammonds, Temporal variation in seismic event rate and b -values from stress corrosion constitutive laws, *Tectonophysics*, *211*, 233-246, 1992.
- Matsu'ura, S. R., Precursory quiescence and recovery of aftershock activities before some large aftershocks, *Bull. Earthquake Res. Inst., Univ. Tokyo*, *61*, 1-65, 1986.
- Meltzner, A. J. and D. J. Wald, Foreshocks and aftershocks of the great 1857 California earthquake, *Bull. Seism. Soc. Am.*, *89*, 1109-1120, 1999.
- Meredith, P. G., I. G. Main and C. Jones, Temporal variation in seismicity during quasi-static and dynamic rock failure, *Tectonophysics*, *175*, 249-268, 1990
- Molchan, G. M. and O. E. Dmitrieva, Aftershock identification - Methods and new approaches, *Geophys. J. Int.*, *109*, 501-516, 1992.
- Molchan, G. M., O. E. Dmitrieva, I. M. Rotvain and J.

- Demey, Statistical analysis of the results of earthquake prediction, based on bursts of aftershocks, *Phys. Earth Planet. Inter.*, 61, 128-139, 1990.
- Morse, P. M. and H. Feshbach *Methods in Theoretical Physics*, McGraw-Hill, New-York, 1953.
- Moyal, J. E., Multiplicative population chains, *Proc Roy. Soc Lond, series A*, 266, 518-526, 1962.
- Ogata, Y., Statistical models for earthquake occurrence and residual analysis for point processes, *J. Am. stat. Assoc.*, 83, 9-27, 1988.
- Ogata, Y., Statistical model for standard seismicity and detection of anomalies by residual analysis, *Tectonophysics*, 169, 159-174, 1989.
- Ogata, Y., Detection of precursory relative quiescence before great earthquakes through a statistical model, *J. Geophys. Res.*, 97, 19845-19871, 1992.
- Ogata, Y., Space-time point process models for earthquake occurrences, *Ann. Inst. stat. Mech.*, 50, 379-402, 1998.
- Ogata, Y., Seismicity analysis through point-process modeling : a review *Pure Appl. Geophys.*, 155, 471-507, 1999.
- Ogata, Y., Increased probability of large earthquakes near aftershock regions with relative quiescence, *J. Geophys. Res.*, 106, 8729-8744, 2001.
- Olver, F. W. J. *Asymptotics and Special Functions* 572 pp., Academic Press, N.Y. and London, 1974.
- Ramselaar, P. A., The mean behavior of the Ogata earthquake process, Master's thesis, Dept of Mathematics, Univ. Utrecht, 1990.
- Ranalli, G., A statistical study of aftershock sequences, *Annali di Geofisica*, 22, 359-397, 1969.
- Reasenberg, P. A., Second-order moment of central California seismicity, *J. Geophys. Res.*, 90, 5479-5495, 1985.
- Reasenberg, P. A., Foreshocks occurrence before large earthquakes, *J. Geophys. Res.*, 104, 4755-4768, 1999.
- Reasenberg, P. A. and L.M. Jones, earthquake hazard after a mainshock in California, *Science*, 243, 1173-1176, 1989.
- Sammis, S. G. and D. Sornette, Positive Feedback, Memory and the Predictability of Earthquakes, *Proc. Nat. Acad. Sci. USA*, 99 SUPP1, 2501-2508, 2002.
- Scholz, C. H., Microfracturing and the inelastic deformation of rocks in compression, *J. Geophys. Res.*, 73, 1417-1432, 1968.
- Simeonova, S. and D. Solakov, Temporal characteristics of some aftershocks sequences in Bulgaria, *Annali di Geofisica*, 42, 821-832, 1999.
- Smith, W. D., The b -value as an earthquake precursor, *Nature*, 289, 136-139, 1981.
- Sornette, D., *Critical Phenomena in Natural Sciences, Chaos, Fractals, Self-organization and Disorder: Concepts and Tools*, Springer Series in Synergetics, Heidelberg, 2000a.
- Sornette, D., Mechanochemistry: an hypothesis for shallow earthquakes, in *Earthquake thermodynamics and phase transformations in the earth's interior*, edited by R. Teisseyre and E. Majewski, International Geophysics Series, Vol. 76, pp. 329-366, Cambridge University Press, 2000b.
- Sornette, A. and D. Sornette, Renormalization of earthquake aftershocks, *Geophys. Res. Lett.*, 26, 1981-1984, 1999.
- Sornette, D. and A. Helmstetter, New Mechanism for Finite-Time-Singularity in Epidemic Models of Rupture, Earthquakes and Starquakes, submitted to *Phys. Rev. Lett.* (e-print at <http://xxx.lanl.gov/abs/cond-mat/0112043>), 2001.
- Steeple, D. W. and D. D. Steeples, Far-field aftershocks of the 1906 earthquake, *Bull. Seism. Soc. Am.*, 86, 921-924, 1996.
- Sykes, L. R. and S. C. Jaumé, Seismic activity on neighboring faults as a long-term precursor to large earthquakes in the San-Francisco Bay region, *Nature*, 348, 595-599, 1990.
- Tajima, F. and H. Kanamori, Global survey of aftershock area expansion patterns, *Phys. Earth Planet. Inter.*, 40, 77-134, 1985.
- Utsu, T., Y. Ogata and S. Matsu'ura, The centenary of the Omori Formula for a decay law of aftershock activity, *J. Phys. Earth*, 43, 1-33, 1995.
- Vere-Jones, D., Statistical theories of crack propagation, *Mathematical Geology*, 9, 455-481, 1977.
- Vere-Jones, D., Robinson, R. and Yang, Remarks on the accelerated moment release model: problems of model formulation, simulation and estimation. *Geophys. Journal. Internat.*, 144, 517-531, 2001.
- Viljoen, H. J., L. L. Lauderback and D. Sornette, Solitary waves and supersonic reaction front in metastable solids, *Phys. Rev. E*, 6502, PT2:U807-U819, 2002.
- Wiemer, S. and K. Katsumata, Spatial variability of seismicity parameters in aftershock zones, *J. Geophys. Res.*, 104, 13,135-13,151, 1999.
- Wyss, M. and R. E. Habermann, Precursory seismic quiescence, *Pure Appl. Geophys.*, 126, 319-322, 1988.
- Yamashita, T. and L. Knopoff, Models of aftershock occurrence, *Geophys. J. Royal Astron. Soc.*, 91, 13-26, 1987.
- Zhang, G., L. Zhu, X. Song, Z. Li, M. Yang, N. Su and X. Chen, Predictions of the 1997 strong earthquakes in Jiashi, Xinjiang, China, *Bull. Seism. Soc. Am.*, 89, 1171-1183, 1999.

Agnès Helmstetter, Laboratoire de Géophysique Interne et Tectonophysique, Observatoire de Grenoble, Université Joseph Fourier, BP 53X, 38041 Grenoble Cedex, France. (e-mail: ahelmste@obs.ujf-grenoble.fr)

Didier Sornette, Laboratoire de Physique de la Matière Condensée, CNRS UMR 6622 Université de Nice-Sophia Antipolis, Parc Valrose, 06108 Nice, France, and Department of Earth and Space Sciences and In-

stitute of Geophysics and Planetary Physics, University of California, Los Angeles, California. (e-mail: sornette@naxos.unice.fr)




Figure 1. Schematic representation of the branching process associated with the ETAS model defined by (1) and (9). In this example, the thickest dashed line is the time arrow associated with the main shock indicated as ‘1’. This main shock triggered five aftershocks denoted ‘11’, ‘12’, ‘13’, ‘14’ and ‘15’ whose magnitudes are proportional to the length of their vertical lines (their position above or below the thickest dashed line is arbitrary and chosen to ensure a better visibility of the diagram). The aftershock ‘11’ triggered three aftershocks denoted ‘111’, ‘112’ and ‘113’. The aftershock ‘12’ triggered four aftershocks denoted ‘121’, ‘122’, ‘123’ and ‘124’. The aftershock ‘13’ triggered a single aftershock denoted ‘131’. The aftershock ‘14’ also triggered a single aftershock denoted ‘141’. The aftershock ‘15’ did not trigger any aftershock. The observable catalog is the superposition of all these events which are projected on the thick dashed line at the bottom of the figure, keeping the thickness as a code for the generation number of each event.




Figure 2. Seismicity rate $N(t)$ in the sub-critical regime with $n = 0.95$. The noisy black line represents the seismicity rate obtained for a synthetic catalog generated using $K = 0.024$, $M = 6.8$, $m_0 = 0$, $c = 0.001$ day, $\alpha = 0.5$, $b = 1.0$ and $\theta = 0.2$, giving the characteristic time is $t^* = 4500$ days. The local Omori law with exponent $p = 1 + \theta = 1.2$ is shown for reference (dotted line). The analytical solution (16) is shown as the thick line. The two dashed lines represents the asymptotic solutions (14) for $t < t^*$ and (15) for $t > t^*$.




Figure 3. Seismicity rate $N(t)$ in the super-critical regime. Same legend as in Figure 2. The synthetic catalog was generated using the same parameters as for Figure 2, except for a lowest b -value of $b = 0.75$ and a smallest mainshock magnitude $M = 6$, leading to a branching number $n = 1.43$ and a characteristic time $t^* = 0.85$ day. The analytical solution (thick line) is calculated from equation (18). The two dashed lines correspond to the approximative analytical solutions (14) and (17).




Figure 4. Seismicity rate $N(t)$ in the case $\theta < 0$ corresponding to a local Omori's law exponent $p < 1$. Same legend as in Figure 2. The synthetic catalog was generated using $K = 0.02$, $M = 7$, $m_0 = 0$, $c = 0.01$ day, $\alpha = 0.5$, $b = 1.0$ and $\theta = -0.1$, giving the characteristic time is $\tau = 10^5$ days. The analytical solution (thick line) is calculated from equation (22). The two dashed lines correspond to the approximative analytical solutions (20) and (21).




Figure 5. Comparison between the three decay laws of aftershock sequences: Omori law with $p = 0.7$ (dashed line), stretched exponential with $q = 1.3$ and $t_0 = 10$ days (thin black line) and our analytical solution in the sub-critical regime (16) for $\theta = q = 1 - p = 0.3$ and $t^* = t_0 = 10$ days (solid gray line). At early times $t \ll t^*$, the three functions are similar and decay as $t^{-0.7}$. At large times, the stretched exponential function and the analytical solution of the ETAS model decay more rapidly than the Omori law. For times up to $t = 10 t^*$, the stretched exponential function is a good approximation of the ETAS model solution, and describes the transition from a power law decay at early times to a faster decay law.




Figure 6. Cumulative aftershock number in the super-critical regime from a synthetic catalog generated using branching ratio $n = 1.27$, $\theta = 0.2$ and $t^* = 4.6$ days. The mainshock magnitude is $M = 7.0$. The thin line is a fit by an Omori law evaluated for time before the occurrence of the first $M \geq 6.0$ aftershock. This fit gives an apparent global p -value of 0.58. Relative seismic quiescence (by comparison with an Omori law) is observed before the occurrence of the $M = 6.0$ aftershock, due to the transition from an Omori law decay with exponent $p = 1 - \theta = 0.8$ for time $t \ll t^*$ to an exponential increase of the seismicity rate for time $t \gg t^*$.

Table 1. ETAS parameters, branching ratio n and characteristic time t^* for the sequences studied by *Ogata* [1989, 1992]. We have computed n and t^* using equations (3) and (12) from the ETAS parameters K , α , c , $p = 1 + \theta$ and μ calculated by *Ogata* [1989, 1992] using a maximum likelihood method. For most sequences, we have assumed $b = 1$ to evaluate n and t^* because b -value is not given in [*Ogata*, 1989, 1992]. Thus, there is a large uncertainty in the n and t^* values in the case where α is close to 1.

Reference	seismicity data	M_0	b	μ day ⁻¹	K	c day	p	α	n	t^* day
<i>Ogata</i> [1989]	Japan, 1895-1980	6.0	1.0	0.005	0.087	0.02	1.0	0.7	Inf	^a
<i>Ogata</i> [1989]	Rat-Island 1963-1982	4.7	1.0	0.0	0.072	0.167	1.35	0.63	1.04	4600
<i>Ogata</i> [1989]	Nagano, 1978-1986	2.5	0.9	0.021	0.008	0.017	0.85	0.94	Inf	^b
<i>Ogata</i> [1989]	Nagano aftershocks, 1986	2.9	1.2	0.0	0.032	0.038	1.14	0.73	0.92	4.10 ⁶
<i>Ogata</i> [1992]	worldwide shallow earthquakes	7.0	1.0	0.019	0.018	0.21	1.03	0.53	1.49	10 ¹⁷
<i>Ogata</i> [1992]	Central Aleutian, 10 years	4.7	1.0	0.008	0.042	0.03	1.13	0.62	1.34	2200
<i>Ogata</i> [1992]	Tohoku, 95 years	6.0	1.0	0.0054	0.98	0.02	1.0	0.70	Inf	^a
<i>Ogata</i> [1992]	Tokachi-Oki aftershocks, 1 year	4.8	1.0	0.14	0.015	0.23	1.28	0.98	4.03	1.5
<i>Ogata</i> [1992]	Niigata aftershocks, 150 days	4.0	1.0	0.075	0.0005	0.15	1.37	1.26	Inf	^b
<i>Ogata</i> [1992]	Niigata aftershocks, 150 days	2.5	1.0	0.47	0.0002	1.10	1.72	1.34	Inf	^b
<i>Ogata</i> [1992]	Izu Islands, 55 years	4.0	1.0	0.0038	0.062	0.012	1.143	0.155	0.96	10 ⁸
<i>Ogata</i> [1992]	Izu Peninsula, 7 years	2.5	1.0	0.022	0.035	0.003	1.35	0.17	0.91	7.3
<i>Ogata</i> [1992]	Off east cost of Izu, 33 days	2.9	1.0	0.59	0.016	0.009	1.73	0.31	1.00	346.
<i>Ogata</i> [1992]	Matsushiro swarm, 20 years	3.9	1.0	0.0006	0.092	0.13	1.14	0.27	1.21	2200
<i>Ogata</i> [1992]	Kanto, 1904-1916	5.4	1.0	0.028	0.010	0.010	1.00	0.62	Inf	^a
<i>Ogata</i> [1992]	Kanto, 1916-1923	5.4	1.0	0.025	0.001	0.010	1.02	1.31	Inf	^b
<i>Ogata</i> [1992]	Hachijo, 1938-1969	5.4	1.0	0.013	0.008	0.004	1.02	0.85	3.0	5.10 ⁶
<i>Ogata</i> [1992]	Hachijo, 1969-1973	5.4	1.0	0.016	0.001	0.013	1.00	1.11	Inf	^a
<i>Ogata</i> [1992]	Tonankai, 1933-1939	5.2	1.0	0.050	0.010	0.065	1.02	0.90	5.28	4.10 ³
<i>Ogata</i> [1992]	Tonankai, 1939-1944	5.2	1.0	0.031	0.009	0.011	1.01	0.83	5.54	10 ⁷
<i>Ogata</i> [1992]	Tokachi, 1926-1945	5.0	1.0	0.047	0.013	0.065	1.32	0.83	0.57	0.40
<i>Ogata</i> [1992]	Tokachi, 1945-1952	5.0	1.0	0.041	5.20	11.6	3.50	1.37	Inf	^b
<i>Ogata</i> [1992]	Tokachi, 1952-1961	5.0	1.0	0.032	0.021	0.059	1.10	0.72	0.99	10 ²²
<i>Ogata</i> [1992]	Tokachi, 1961-1968	5.0	1.0	0.014	0.014	0.005	0.86	0.43	Inf	7.10 ^{5c}

^a t^* cannot be evaluated because $p = 1$

^b t^* cannot be evaluated because $\alpha < b$

^c τ is given instead of t^* because $\theta < 0$